



[urn:lsid:zoobank.org:pub:A3A6CDB8-BDC8-4CD4-A409-77508FC11530](https://zoobank.org/pub:A3A6CDB8-BDC8-4CD4-A409-77508FC11530)

Redescription of *Proformica nasuta* (Nylander, 1856) (Hymenoptera, Formicidae) using an integrative approach

Christophe GALKOWSKI¹, Claude LEBAS², Philippe WEGNEZ³,
Alain LENOIR⁴ & Rumsais BLATRIX^{5,*}

^{1,2,3,4,5} AntArea (www.antarea.fr), Association for the Study and
Mapping of Ants from Metropolitan France.

¹ 104 Route de Mounic, 33160 Saint-Aubin-de-Medoc, France.

² 2 Impasse del Ribas, 66680 Canohès, France.

³ Walbru (www.fourmiswalbru.com), Belgian Association for the Inventory of Ant Species in
Wallonia and Brussels, and Rue de la Grotte 23, 4651 Herve, Belgium.

⁴ IRBI, Institut de Recherche sur la Biologie de l’Insecte, UMR CNRS 7261,
Université François Rabelais, Faculté des Sciences, Parc Grandmont, 37200 Tours, France.

⁵ CEFE UMR 5175, CNRS – Université de Montpellier – Université Paul Valéry
Montpellier – EPHE, 1919 Route de Mende, 34293 Montpellier Cedex 5, France.

* Corresponding author: rumsais.blatrix@cefe.cnrs.fr

¹ E-mail: chris.gal@wanadoo.fr

² E-mail: cllebas@free.fr

³ E-mail: wegnez.phil@gmail.com

⁴ E-mail: alain.lenoir@univ-tours.fr

¹ [urn:lsid:zoobank.org:author:C480F514-D567-4787-AF55-2BE00FB79706](https://zoobank.org/author:C480F514-D567-4787-AF55-2BE00FB79706)

² [urn:lsid:zoobank.org:author:99DB1C68-3E40-48A7-A665-8BDAF6D8D1C3](https://zoobank.org/author:99DB1C68-3E40-48A7-A665-8BDAF6D8D1C3)

³ [urn:lsid:zoobank.org:author:44194EA4-001C-4CE0-9567-F9977E103B40](https://zoobank.org/author:44194EA4-001C-4CE0-9567-F9977E103B40)

⁴ [urn:lsid:zoobank.org:author:0B03EB0F-06EB-413C-A91E-B57CBB22C623](https://zoobank.org/author:0B03EB0F-06EB-413C-A91E-B57CBB22C623)

⁵ [urn:lsid:zoobank.org:author:AFF6B4B9-C054-43A2-8095-84F0FB7B6EE9](https://zoobank.org/author:AFF6B4B9-C054-43A2-8095-84F0FB7B6EE9)

Abstract. The taxonomy of the Palaearctic ant genus *Proformica* Ruzsky, 1902 is confused and in need of revision. The type specimen for *P. nasuta* (Nylander, 1856), the type species of the genus, was from Beaucaire, southern France, and is presumably lost. Based on extensive sampling of *Proformica* nests in southern France, including the type locality, we show that the concept of *P. nasuta* has been erroneous for more than a century. We integrate information from the morphology of workers and sexual castes, DNA markers, and cuticular hydrocarbons to re-define species in southern France. This allowed us to provide a new, accurate description of *P. nasuta* and designate a neotype, as well as reference individuals for all castes. In addition, we propose a name, *P. longipilosa* sp. nov., for a species that since the end of the 19th century has mistakenly been included in *P. nasuta*.

Keywords. Ants, *Proformica longipilosa* sp. nov., Western Palaearctic, molecular markers, cuticular hydrocarbons.

Galkowski C., Lebas C., Wegnez P., Lenoir A. & Blatrix R. 2017. Redescription of *Proformica nasuta* (Nylander, 1856) (Hymenoptera, Formicidae) using an integrative approach. *European Journal of Taxonomy* 290: 1–40. <https://doi.org/10.5852/ejt.2017.290>

Introduction

The ant genus *Proformica* Ruzsky, 1902 is composed of 25 species (Bolton 2014) restricted to dry and open environments such as steppes, mountain meadows and Mediterranean seashores (Agosti 1994). It is endemic to the Palaearctic region, with a disjunct distribution. A first area extends from eastern Europe to eastern Asia and contains most of the species, and a second area, much more limited in species number and distribution, occurs at the southwestern tip of Europe (Portugal, Spain and southern France). This distribution is somewhat reminiscent of that of the meadow and steppe vipers, the *Vipera ursinii* species complex, which is composed of taxa restricted to steppe-like ecosystems. Asia and Europe show distinct viper taxa that diverged in the early Pliocene, about 4 Mya (Zinenko *et al.* 2015). The genus *Proformica* may have experienced the same biogeographic history as these vipers and several other organisms inhabiting steppe-like environments (Ruano *et al.* 2011; Sanllorente *et al.* 2015). Only one taxon, *P. nasuta* (Nylander, 1856), is reported to occur in both Asian and western European areas.

The taxonomy of the genus *Proformica* is complicated and in need of revision. The situation is particularly complex in the eastern area, with currently 23 species reported. In Western Europe, two distinct zoogeographical areas can be distinguished, the Iberian Peninsula and southern France, which are separated by a barrier formed by the Pyrenees mountain range. Three described species are currently recorded for the Iberian Peninsula (Collingwood 1976), but at least six forms are recognized by ant taxonomists (Xavier Espadaler, Barcelona, pers. comm.) and substantial morphological variation within each form makes species delimitation difficult. In contrast, only one described species, *P. nasuta*, has been recorded for southern France (*P. ferreri* Bondroit, 1918 may also be present in the French part of the Pyrenees).

Proformica nasuta is the type species for the genus *Proformica* and was described from Beaucaire, France. The concept of this species is unclear. For instance, variation in the number of erect hairs on the mesosoma, a character commonly used in the taxonomy of *Proformica*, has been interpreted either as mere intraspecific variation (Espadaler & Cagniant 1987), or as an indication that the name *P. nasuta* actually covers two taxa (Santschi 1925; Collingwood & Yarrow 1969). Populations of species of *Proformica* are small, inconspicuous and patchily distributed, and the species are often considered rare. As a consequence, the genus is poorly represented in institutional collections and most taxonomic work is based on few specimens, rendering the accurate perception of intra-specific variation difficult. Moreover, the type specimen of *P. nasuta* has not been located. Having not been found in the most likely candidate collections and not explicitly referred to in the literature, it is presumably lost. As *P. nasuta* is the oldest name in the genus, designation of a neotype and a precise redefinition of this taxon are indispensable before further taxonomic work on this genus can be undertaken. For this purpose, we analysed a sample, unprecedented in its size and geographic extent, of *Proformica* nests in southern France using an integrative taxonomy approach based on morphological data from workers and sexuals, DNA sequences and cuticular hydrocarbons. Southern France was the best location for this investigation as it encompasses the type locality for *P. nasuta* and harbours no other known *Proformica* species. Combining the results of these different characters can increase our ability to provide valid decisions about species delimitations (Schlick-Steiner *et al.* 2010). Although some of these kinds of data are less relevant than others for the descriptive taxonomy of a particular species group, incongruences between results based on different kinds of data can provide information on the biology of the group studied and insights into ongoing ecological and evolutionary processes.

Material and methods

Sampling

We sampled 110 nests of *Proformica* at 22 localities from across southern France (Fig. 1), including four nests at the type locality of *P. nasuta* (Beaucaire) and seven nests less than 10 km from that site. Collection details for each nest are provided in Appendix 1. We give here a more detailed description of the collection localities. Fourteen sites (Aurons, Beaucaire, Bonnieux, Collias, Plaine de la Crau, Grospierres, Jonquières, Montpellier, Orange, Pompignan, Sauteyrargues, Sisteron, Tarascon, Vinsobre) were found in lowland Mediterranean garrigue or sun-exposed grassland. Nests were usually found in areas where soil had been heavily trampled. Four sites (Grand Luberon, Montagne de Lure, Sainte-Baume, Sumène) were on rocky summits of medium-sized calcareous mountains with little vegetation cover. Finally, four sites (Plateau de Calern, Plateau de Caussols, Gréolières, Mont Ventoux) were on mountain slopes or plateaus in the foothills of the Alps, also bearing little vegetation cover.

Morphological investigation

Most of the morphological characters used were introduced by Seifert (2007).

Eight morphological characters were measured on 321 worker individuals originating from 97 nests collected at 21 localities (Appendix 1). Colonies are monodomous, so each nest represents one colony. Mean values were calculated for worker individuals collected from each nest (range 1–8, mean 3.3 workers per nest).




Fig. 1. Distribution of the 22 localities where *Proformica* nest samples were collected. The star indicates the type locality of *Proformica nasuta* (Nylander, 1865) (i.e., Beaucaire).

Morphometric characters measured:

- CW = Maximum cephalic width (including eyes) (in μm)
GHL = Length of the longest erect seta on the gaster (in μm)
nCH = Unilateral number of erect setae on the vertex
nCU = Unilateral number of erect setae on the underside of the head
nG = Unilateral number of erect setae on the gaster
nSc = Unilateral number of erect setae on the petiole
nTx = Unilateral number of erect setae on the mesosoma
PDG = Pubescence distance on dorsum of the first gaster tergite, computed as L/N , where N is the number of pubescent (appressed) hairs crossing a transverse line of length L ; here we used a line of length $340 \mu\text{m}$

Ten queens from six localities (Beaucaire, Orange, Sainte-Baume, Pompignan, Tarascon, Mont Ventoux) were measured for 14 characters:

- ML = Mesosoma length (in μm)
MW = Mesosoma width (in μm)
nMes = Unilateral number of erect setae on the scutum and scutellum, or on the mesonotum if scutum and scutellum were not discernible
nPn = Unilateral number of erect setae on the pronotum
nPP = Unilateral number of erect setae on the propodeum
ScW = Width of petiole scale (in μm)
SL = Scape length (in μm)
CW, nCH, nCU, nSc, nG, PDG and GHL: see worker morphology.

Measurements of workers and queens are given below (Appendices 2–3).

We collected a total of 11 males from three localities (Plaine de la Crau, Pompignan and Tarascon) and examined males from Sainte-Baume that had been collected by F. Bernard (MNHN) in July 1974. Males were only described qualitatively.

Acronyms of depositories:

- AT = Alberto Tinaut Collection, University of Granada, Granada, Spain
IRSNB-BC = Bondroit Collection, Institut Royal des Sciences Naturelles de Belgique, Brussels, Belgium (contact: Wouter Dekoninck)
LB = Personal collection of Lech Borowiec, University of Wroclaw, Wroclaw, Poland
MCSN-EC = Emery Collection, Museo Civico di Storia Naturale, Genova, Italy (contact: Roberto Poggi)
MCZL-KC = Kutter Collection, Musée Cantonal de Zoologie, Lausanne, Switzerland (contact: Anne Freitag)
MHNG = Muséum d'Histoire Naturelle de Genève, Geneva, Switzerland (contact: Bernhard Merz)
MHNG-FC = Forel Collection, Muséum d'Histoire Naturelle de Genève, Geneva, Switzerland (contact: Bernhard Merz)
MNHN = Muséum national d'Histoire naturelle, Paris, France
MNHN-BAC = Collections from Bernard, André and others, Muséum national d'Histoire naturelle, Paris, France (contact: Agnès Touret-Alby and Quentin Rome)
NMB-SC = Santschi Collection, Naturhistorisches Museum, Basel, Switzerland (contact: Isabelle Zürcher-Pfander)

SMNH	=	Senckenberg Museum of Natural History, Görlitz, Germany
XE	=	Personal collection of Xavier Espadaler (Autonomous University of Barcelona, Barcelona, Spain)
ZISP	=	Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia

Molecular phylogenetic analysis

DNA was extracted from 1–2 individuals from each of 98 nests distributed in 20 localities (Appendix 1), using the REExtract-N-Amp PCR Kit (Sigma–Aldrich, St. Louis, MO). Amplification by polymerase chain reaction (PCR) was performed using either the Goldstar Red DNA Polymerase (Eurogentec), the Qiagen Multiplex kit (Qiagen, Venlo, Netherlands) or the Sigma REExtract-N-Amp PCR Ready Mix (Sigma–Aldrich). Conditions for PCR first followed manufacturer’s instructions, and were then adjusted if amplification failed. Sanger sequencing of PCR products was performed at the Genoscope (Evry, France).

We amplified four DNA markers, two mitochondrial, two nuclear: (i) COI (~600 bp), coding for part of the cytochrome c oxidase subunit 1, (ii) Cytb (~700 bp), corresponding to the end of the sequence coding for NADH dehydrogenase subunit 6 and part of cytochrome b, (iii) 28S (~600 bp), coding for part of the large ribosomal subunit, and (iv) LW Rh (~550 bp), coding for part of the long-wavelength rhodopsin. COI was amplified for 45 *Proformica* individuals (GenBank accession numbers: KU749600–KU749637 and KU749641–KU749654) using two sets of primers covering the same region: either LepF1 (5'-ATTCAACCAATCATAAAGATAT-3') and LepR1 (5'-TAAACTTCTGGATGTCCAAAAA-3') (Hebert *et al.* 2004), or CI13 (5'-ATAATTTTTTTTATAGTTATAACC-3') and CI14 (5'-ATTTCTTTTTTTT-CCTCTTTC-3') (Hasegawa *et al.* 2002). For some individuals we used the two different primer pairs and compared the sequences obtained for the same individual. For each of four individuals, two highly divergent copies of COI were sequenced. To detect sequences that might come from accidental amplification of numts (copies of mitochondrial DNA transferred into the nuclear genome), we searched for the presence of premature stop codons in the amino-acid sequences. Three sequences (belonging to two individuals) had one premature stop codon. Based on the distribution of these sequences and the divergent copies from the same individual in the COI phylogeny, we identified a clade of putative numts. We amplified the Cytb marker using primers Cytb-FeF (5'-CAGTTTAATTTCTAATGAACAAAC-3') and Cytb-FeR (5'-GGATCTCTAAAAATATATGGG-3') (Liautard & Keller 2001) for a subset of *Proformica* individuals, and we used these sequences to design internal primers more specific to *Proformica* in order to increase amplification success: cytbPf (5'-CCTTTTAATAATRTYACTATTGC-3') and cytbPr (5'-TATAARTTTCTATTAATTCCAAG-3'). In total we amplified the Cytb marker in 103 individuals of *Proformica* (GenBank accession numbers: KU749655–KU749737 and KU749739–KU749758). The 28S marker was amplified for 31 individuals of *Proformica* (GenBank accession numbers: KU749759–KU749783 and KU749788–KU749793) using primers D2B (5'-GTCGGGTTGCTTGAGAGTGC-3') (Saux *et al.* 2004) and D2R (5'-TTGGTCCGTGTTTCAAGACGGG-3') (Belshaw & Quicke 1997). The LW Rh marker was amplified for 31 individuals of *Proformica* (GenBank accession numbers: KU749794–KU749818 and KU749823–KU749828) using primers LR143F (5'-GACAAAGTKCCACCRGARATGCT-3') and LR639ER (5'-YTTACCGRTTCCATCCRAACA-3') (Ward & Downie 2005). DNA sequences were aligned with MUSCLE (Edgar 2004). Alignments were inspected visually and edited manually using MEGA5 (Tamura *et al.* 2011) when they could be improved. Alignment of the intergenic region in the Cytb marker was ambiguous, and thus removed from the analysis.

As outgroup we used species for which we obtained new sequences (Appendix 1): *Bajcaridris theryi* (Santschi, 1936) (GenBank accession numbers, 28S: KU749786, LW Rh: KU749821), *Cataglyphis cursor* (Fonscolombe, 1846) (GenBank accession numbers, COI: KU749638 and KU749640, Cytb: KU749738, 28S: KU749787, LW Rh: KU749822), *Cataglyphis mauritanica* (Emery, 1906) (GenBank

accession numbers, COI: KU749639, 28S: KU749785, LW Rh: KU749820), *Formica cunicularia* Latreille, 1798 (GenBank accession numbers, 28S: KU749784, LW Rh: KU749819), and species for which sequences were retrieved from GenBank (Hasegawa *et al.* 2002; Goropashnaya *et al.* 2004, 2007, 2012; Ward & Downie 2005; Moreau *et al.* 2006): *Cataglyphis iberica* (Emery, 1906) (COI: DQ353343), *Formica cunicularia* (COI: AB010926), *Formica exsecta* Nylander, 1846 (COI: AB010927, Cytb: JX170868), *Formica pratensis* Retzius, 1783 (Cytb: AY584233), *Formica rufibarbis* Fabricius, 1793 (Cytb: JX170889), *Formica truncorum* Fabricius, 1804 (COI: AB010929), *Formica uralensis* Ruzsky, 1895 (Cytb: JX170879), *Formica wheeleri* Creighton, 1935 (28S: DQ353556, LW Rh: DQ353149).

A partition scheme was defined with PartitionFinder (Lanfear *et al.* 2012) for each phylogenetic analysis, using the Bayesian Information Criteria for nucleotide substitution model selection. Prior data blocks were defined by marker and codon position. Three separate phylogenetic reconstructions were performed using both maximum likelihood and Bayesian inference algorithms: one for COI (to highlight the position of the clade of putative numts), one for Cytb (which includes the largest number of individuals), and one for the concatenated nuclear markers (28S + LW Rh) (because nuclear and mitochondrial markers might tell different stories).

Maximum likelihood phylogenies were constructed with RAxML (Stamatakis *et al.* 2008) on the web server at vital IT, Switzerland (<http://embnet.vital-it.ch/raxml-bb/>), using the GAMMA model of rate heterogeneity. Node support was estimated by generating 100 trees by bootstrapping. Bayesian inference phylogenies were constructed with MrBayes 3.2 (Ronquist *et al.* 2012). For the COI phylogeny we used the substitution models SYM+G, F81 and GTR+G for the first, second and third codon position, respectively. For Cytb we used HKY+G, HKY+I and GTR+I+G for the first, second and third codon position respectively. For the concatenated nuclear genes we used K80 for the first codon position of LW Rh, and K80+I for 28S and the second and third codon positions of LW Rh. Each analysis consisted of two runs of four Markov chains run for 10 million generations. Parameters were unlinked for all partitions. A standard deviation of split frequencies of less than 0.01 between two independent runs was reached after less than 2.4 million generations. A burn-in fraction of the first 25 % of the trees was discarded.

Cuticular hydrocarbons

Colonies from nine localities (Bonnieux, Plaine de la Crau, Montpellier, Grand Luberon, Montagne de Lure, Sainte-Baume, Pompignan, Sisteron, Mont Ventoux) were used for analysis of cuticular hydrocarbons. Using forceps, we gathered three to five workers from each colony and put them into glass vials containing 1 ml of hexane. The containers were stored in a freezer at -20°C until chemical analysis. For chemical analysis, the ants were retrieved from the vials and the solvent evaporated. The extract was re-dissolved in 10 µl of hexane. Two µl of each extract were injected into a Perkin-Meyer GC-MS functioning at 70eV with a source temperature of 230°C and equipped with a ZB-5HT column (30 ml × 0.25 mm ID × 0.252 µm df; 5% phenyl- 95% dimethylpolysiloxane). The temperature program was 2 min at 150°C, and then 5°C/min until 320°C, and a 5 min hold at 320°C (total 41 min). Substances were identified using standard alkanes, library data and Kovats retention indices. For the comparisons, we calculated the percentage of each hydrocarbon from the total hydrocarbon content in each ant sample. The data were analysed using Principal Component Analysis. We chose not to transform the data since transformation introduces additional background noise into the data when numerous zero values are present; these have to be replaced to make transformation possible when comparing species. Indeed, reanalysis of the data after transformation (following the procedure of Reymont 1989) gave similar results, but with slightly less efficient separation of groups (Oppelt *et al.* 2008). Analyses were made with the Statistica software.

We also performed chromatograms of cuticular hydrocarbons for two species used as outgroups: *Proformica longiseta* Collingwood, 1978 from Sierra Nevada (Spain) and *Cataglyphis cursor* from Aix-en-Provence (France). Lists of cuticular hydrocarbons known for these species have been published in Errard *et al.* (2006) and Nowbahari *et al.* (1990), respectively, but without quantification.

Nest census and queen reproductive status

Six nests were excavated in July 2011 and the ants counted. Six queens from two nests were dissected to assess their reproductive status. In addition, one apterous queen was obtained by rearing pupae from Sainte-Baume and was dissected to confirm its queen status. Several workers of various sizes were also dissected.

Results

Morphology and altitudinal distribution

Two groups of nests were separated by combining two morphological characters, GHL and PDG for the workers, and GHL and nG for the queens (Fig. 2). The two groups were distinct for both characters, independently of CW, a proxy for size (Fig. 2). One of the groups, coloured in orange in the figures and hereafter denominated as the orange taxon, encompasses the type locality of *Proformica nasuta* (Fig. 1). The other group is coloured in blue in the figures and is hereafter denominated as the blue taxon. Workers of the orange group present denser pubescence, shorter erect hairs on the gaster and fewer erect hairs (or none) on the mesosoma (PDG < 29, GHL < 77 μ m, nest means, nTx \times 2 \pm SD = 9.5 \pm 10.4) than those of the blue group (PDG > 33, GHL > 85 μ m, nest means, nTx \times 2 \pm SD = 46.8 \pm 16.7).

Further, queens of the blue taxon were all winged or showed wing sclerites, while all queens of the orange taxon were ergatoid. We did not find males of the blue taxon in the field and we could not locate specimens in museum collections. Males of the orange taxon have dense and long hairs on the head, mesosoma and the anterior face of the first gaster segment.

For both the orange taxon and the blue taxon, altitudinal distribution of the nests was bimodal (Fig. 3). Most nests of the orange taxon were found below 200 m, but those from Sainte-Baume, Sumène, Grand Luberon and Montagne de Lure were found above 800 m, at the tops of medium-sized mountains. In contrast, most nests of the blue taxon were found above 1000 m on plateaus and mountains, but those from Orange, Sisteron and Vinsobres were found lower, below 600 m. Interestingly, within each of these taxa, GHL and PDG are highest for workers from the mountain localities (except for Sumène) (Fig. 2).

Molecular phylogenetic analysis

Maximum likelihood and Bayesian inference produced very similar phylogenies, so we chose to present only Bayesian inference phylogenies. The clade of putative numts in the COI phylogeny is delimited in red in Fig. 4A. The two mitochondrial markers yielded similar topologies (Fig. 4A–B), showing two main clades which corresponded approximately to the two taxa defined in the morphological analysis. Mismatch between morphotypes and clades was observed for some specimens. The nuclear markers showed very little variation. As a consequence, the resulting tree is poorly resolved (Fig. 4C).

Cuticular hydrocarbons

Identification of compounds and examples of chromatograms are included below (Appendices 4–5). Chromatograms showed very different profiles. Most hydrocarbons of the blue taxon are shorter (< C28) than those of the orange taxon (> C28). Two forms can be distinguished in the orange taxon: the lowland localities, in which individuals have the longest hydrocarbons (> C31), and the mountain localities (Montagne de Lure and Grand Luberon), where the hydrocarbons are of an intermediate length




Fig. 2. Distribution of morphological characters of *Proformica* workers (left) and queens (right) collected in southern France. Top and middle graphics represent the regression of gaster hair length (GHL), pubescence distance on the gaster (PDG) and unilateral number of hairs on the gaster (nG) against cephalic width (CW), a proxy for size. The graphics at the bottom represent a combination of two morphological characters that highlights two distinct groups of nests. These two groups are coloured in blue and orange respectively. The dots with a black cross correspond to nests from mountain and lowland localities for the orange and blue groups, respectively.

(Appendix 5). The PCA distinguishes the two outgroups, *Proformica longiseta* and *Cataglyphis cursor*, from the *Proformica* samples from southern France (Fig. 5). The blue and orange taxa are segregated along the first axis of the PCA. In addition, the strongest differentiation occurs within the orange taxon, between a group formed by the two mountain localities (Montagne de Lure and Grand Luberon) and the others.

Queen reproductive status, nest census

Excavation of nests of the two taxa revealed the same general structure: the entrance opens directly at the ground surface, sometimes under a small stone; a vertical gallery of 10–20 cm leads to a small chamber where males can be found when present; then, the gallery goes down obliquely and reaches a final chamber, about 50 cm below ground level, where queens are present. Secondary galleries, lateral (perpendicular) to the principal one, may be present and lead to chambers. The content of nests is presented in Table 1. Repletes, i.e. workers with inflated gaster serving as stores of liquid food, were




Fig. 3. Altitudinal distribution of 110 nests of *Proformica* from southern France belonging to two morphological groups.

Taxonomy

Class Hexapoda Blainville, 1816
 Order Hymenoptera Linnaeus, 1758
 Superfamily Vespoidea Latreille, 1802
 Family Formicidae Latreille, 1809
 Subfamily Formicinae Latreille, 1809

***Proformica* Ruzsky, 1902**

Nylander (1856) described *P. nasuta*, the type species of the genus, from Beaucaire. Our analyses assigned workers from the type locality and from two other localities within a radius of 10 km (Jonquières and Tarascon) to the orange taxon. They lack erect hairs on the mesosoma, agreeing with the description of *Proformica nasuta* by Nylander as “nuda”. Although the type is presumably lost (as it could not be found in the following collections: Nylander (Helsinki) (Radchenko 2007), Forel (Geneva), Emery (Genoa), Bondroit (Brussels) and Santschi (Basel)), we are confident that the nest samples we collected in Beaucaire and in the surrounding area correspond to the species described by Nylander. Below we provide a redescription of *P. nasuta* (the orange taxon), and the description of a new species, *P. longipilosa* sp. nov. (the blue taxon).




Fig. 5. Principal Component Analysis of *Proformica* localities based on relative proportions of cuticular hydrocarbons identified by gas chromatography and mass spectrometry. Colours match the groups defined in the morphological analysis.

Table 1. Content of excavated nests of *Proformica Ruzsky, 1902*. * = a large part of the nest could not be collected. The numbers of individuals are underestimates.

Colony	Taxon colour code	Species	No. of queens	Queen status	No. of males	No. of repletes	No. of other workers	Total no. of workers
Beaucaire 1	orange	<i>P. nasuta</i>	2	apterous	0	11	65	76
Pompignan 10	orange	<i>P. nasuta</i>	16	apterous	0	87	515	602
Tarascon 1	orange	<i>P. nasuta</i>	1	apterous	5	17	121	138
Orange 1	blue	<i>P. longipilosa</i>	7	wing sclerites	0	9	226	235
Mont Ventoux 10*	blue	<i>P. longipilosa</i>	1	wing sclerites	0	4	339	343
Mont Ventoux 14*	blue	<i>P. longipilosa</i>	5	wing sclerites	0	79	91	170

Table 2. Ovary status of seven queens of *Proformica Ruzsky, 1902*. * = queen emerged after rearing pupae.

Colony	Queen type	Taxon colour code	Species	Spermatheca	Mature oocytes	Yellow bodies	No. of ovarioles per ovary
Pompignan 10	apterous	orange	<i>P. nasuta</i>	full	many	many	~10
Pompignan 10	apterous	orange	<i>P. nasuta</i>	full	many	many	~10
Pompignan 10	apterous	orange	<i>P. nasuta</i>	full	many	many	~10
Sainte-Baume*	apterous	orange	<i>P. nasuta</i>	empty	absent	absent	~10
Orange 1	dealate	blue	<i>P. longipilosa</i>	full	many	many	> 15
Orange 1	dealate	blue	<i>P. longipilosa</i>	full	many	many	> 15
Orange 1	dealate	blue	<i>P. longipilosa</i>	full	many	many	> 15

Redescription of *P. nasuta* (Nylander, 1856) and designation of the neotype

As the type specimen of *P. nasuta* is presumably lost, we propose fixation of a neotype from a nest sample collected in Beaucaire, France, *terra typica* of the species, and matching Nylander's concept of *P. nasuta*. The original description (Nylander 1856: 66) is based on a small worker ("Long. 3 – 3.5 mm") with elongated head ("...*facies producta antice visa subrectangularis...*"). This feature is found exclusively in minor workers. Therefore, a minor worker was selected from Beaucaire, France (colony Beaucaire 1) and designated as the neotype. The neotype is deposited as MNHN-1598 with the labels "FRA, N43.83544 E4.61828, Beaucaire, 9 juillet 2011, leg. R. Blatrix & C. Lebas" and "Néotype *Proformica nasuta* (Nylander, 1856), des. Galkowski, Lebas, Wegnez, Lenoir & Blatrix, 2016". In case of loss or destruction of this specimen, a replacement neotype can be designated from a series of ten other minor workers collected from the same nest and deposited at the MNHN. Other workers from the same nest are deposited at the following collections: AT (no. 15557), LB, SMNH, XE, ZISP and the collections of the authors. A queen from the same nest and a male from colony Tarascon 1 (a few kilometers away from the type locality) are deposited at MNHN.

***Proformica nasuta* (Nylander, 1856)**
Figs 6–8

Formica nasuta Nylander, 1856: 66.

Formica (Proformica) nasuta var. *depilis* Santschi, 1925: 353.

Material examined**Museum material**

FRANCE: MHNG: Charleval, Bouches-du-Rhône, leg. E. della Santa, labelled *P. nasuta*, 11 Jul. 1987: 4 workers, 25 Jul. 1988: 3 workers. – NMB-SC: Carrière des Anglais, Vaucluse: 2 workers, types of *Formica (Proformica) nasuta* var. *depilis*. – MNHN-BAC: Sainte-Baume, ~90 workers (12 pins), 3 ♂♂, one of which labelled "Type *phoenica*" [*Proformica phoenica* is a *nomen nudum*, as it was never described by Bernard. *Proformica* ants at this locality form a mountain isolate which most probably derived from lowland populations of *P. nasuta*; see Discussion]. – IRSNB-BC: 2 ♂♂, without locality.

New material

FRANCE: All in personal collections of CF and RB: Aurons, 4 workers, 43.66° N, 5.15° E, Jun. 2006, leg. V. Bouchet; Beaucaire, 14 workers, 2 queens, 43.8354° N, 4.6187° E, 9 Jul. 2011, leg. R. Blatrix and C. Lebas; Bonnieux, 8 workers, 43.8625° N, 5.3069° E, 1 Oct. 2011, leg. C. Lebas; Collias, 5 workers, 43.9477° N, 4.4623° E, 12 Jun. 2010, leg. R. Blatrix; Plaine de la Crau, Saint-Martin-de-Crau, 18 workers, 1 ♂, 43.5833° N, 4.8333° E, Jun. and Aug. 2011, leg. C. Lebas; Grospierres, 30 workers, 44.4116° N, 4.2713° E, 6 Jul. 2012, leg. T. Colin; Jonquières-Saint-Vincent, 7 workers, 43.8314° N, 4.5765° E, 11 May 2011, leg. R. Blatrix; Montpellier, 3 workers, 43.6292° N, 3.8907° E, 8 Mar. 2012, leg. R. Blatrix; Pompignan, 3 workers, 5 queens, 2 ♂♂, 43.8979° N, 3.8252° E, May 2010 and 10 Jul. 2011, leg. R. Blatrix and P. Wegnez; Sauteyrargues, 4 workers, 43.8275° N, 3.9192° E, 28 May 2011, leg. R. Blatrix; Sumène, 6 workers, 43.9904° N, 3.7714° E, 9 Apr. 2014, leg. R. Blatrix; Tarascon, 12 workers, 1 queen, 3 ♂♂, 43.8421° N, 4.7382° E, 9 Jul. 2011, leg. R. Blatrix and C. Lebas.

Description**Minor worker** (n = 89)

For each character extreme values and the mean are given in brackets:

CW 540–900 µm; nCH 0–2 (0.06); nCU 0–3 (0.18); nTx 0–6 (0.57); nSc 0–2 (0.13); nG 0–5 (1); PDG 9–28 (13.2); GHL 35–75 µm (45.2); GHL/CW 0.040–0.102 (0.064).




Fig. 6. *Proformica nasuta* Nylander, 1856, neotype, minor worker from colony Beaucaire 1, France. A. Lateral view. B. Dorsal view. C. Head in full face view. D. Gaster in dorsal view. Scale bars = 1 mm. Automontage: Claude Lebas.

Media and major workers (n = 25)

CW 900–1240 μm ; nCH 0–1 (0.13); nCU 0–4 (1.13); nTx 0–7 (1.6); nSc 0–4 (0.27); nG 0–7 (2.3); PDG 10–18 (14.2); GHL 35–90 μm (60.2); GHL/CW 0.039–0.091 (0.054).

Body uniformly dark brown to black, appendices and mandibles lighter. Erect hairs rare or absent, short when present (GHL/CW < 0.11). Dense pubescence on dorsal surface of first and second gaster tergites (PDG < 29). Profile of mesosoma sinuous. Petiolar scale erect, thick, slightly notched at summit in large workers. Head of minor workers clearly elongate, rectangular (CL/CW > 1.3). Head of media and major workers less elongate (CL/CW 1.1–1.3), a bit shiny toward occiput, faintly sculptured in anterior part. Clypeus finely striate longitudinally. Mandible with five teeth of increasing size from base to apex.

Queen (n = 8)

CW 1500–1690 μm (1600); CL 1530–1660 μm (1590); SL 1230–1270 μm (1250); nCH 0–1 (0.2); nCU 2–6 (3.2); nTx 2–19 (11.8); nSc 0–2 (1.2); nG 0–3 (1.6); PDG 9–14 (11.2); GHL 70–90 μm




Fig. 7. *Proformica nasuta* Nylander, 1856, queen from colony Beaucaire 1, France. **A.** Lateral view. **B.** Dorsal view. **C.** Head in full face view. Scale bars = 1 mm. Automontage: Claude Lebas.

(77.5); MW 1020–1240 μm (1150); ML 2040–2310 μm (2200); ScW 620–860 μm (710);
GHL/CW 0.042–0.056 (0.048).

Only ergatoid queens collected. Body brown with a wide orange spot on mesosoma on some specimens.
Few and relatively short erect hairs (nG < 10, GHL < 90 μm). Pubescence very dense on the dorsal




Fig. 8. *Proformica nasuta* Nylander, 1856, male from colony Tarascon 1, France. **A.** Lateral view. **B.** Head in full face view. **C.** Genitalia in dorsal view. Scale bars = 1 mm. Automontage: Claude Lebas.

surface of first and second gaster tergites. Profile of mesosoma similar to that of worker, but mesonotum more domed and propodeum very high. Tegulae absent. Petiolar scale high and wide, distinctly notched at summit. Head as long as wide. Same sculpture as in worker.

Male (n = 6)

CW 1340–1380 μm (1360); CL 1100–1170 μm (1140); SL 1220–1330 μm (1280); MW 1700–1750 μm (1730); ML 2720–2970 μm (2830); ScW 560–590 μm (580).

Head, mesosoma and scale dark brown, appendices and sometimes gaster lighter. Erect hairs numerous and dense on entire head, mesosoma and petiolar scale, becoming rare on gaster, where restricted to anterior face of first tergite, angled. Pubescence almost lacking on dorsal surface of gaster tergites. Eyes and ocelli prominent. Mandibles reduced, without teeth. Many erect black hairs on extensor profile of anterior and median femurs. Maximal length of these hairs shorter than width of femur. Rare erect white hairs on tibiae. Wings well developed, yellowish.

Distribution

Southern France (Fig. 1).

Remarks

We have not examined the type specimens of the Asian varieties of this species, *P. nasuta metalica* Kuznetsov-Ugamsky, 1923 and *P. nasuta syrdariana* Santschi, 1928, described from Kazakhstan, nor the type specimen of the taxon *Formica aerea* (Roger, 1859), which was described based on a single minor worker collected in Greece and later synonymized with *P. nasuta* by Emery (1925). Although we decided not to change the status of these eastern taxa until a thorough revision of the eastern *Proformica* is undertaken, we believe it is very unlikely that they will be conspecific with *P. nasuta*.

Proformica longipilosa sp. nov.

[urn:lsid:zoobank.org:act:6EF784CD-C8B2-4474-840E-04AFE460C0CE](https://doi.org/10.21203/rs.3.rs-1234567)

Figs 9–10

Diagnosis

Workers varying greatly in size, the smallest having a strongly elongated head. Body black. Pubescence sparse and sculpture of tegument weak, giving a shiny aspect. Hairs on mesosoma and gaster very long.

Etymology

The epithet of this species refers to the long erect hairs on the mesosoma and gaster of workers.

Material examined

Holotype

FRANCE: minor worker from colony Mont Ventoux 10 of this study, 44.15261° N, 5.32081° E, alt. 1442 m, 14 Jul. 2011, coll. R. Blatrix (MNHN 1598).

Paratypes

FRANCE: Same collection data (including nest) as holotype: 10 workers, (MNHN 1598); other workers in AT (15556), LB, SMNH, XE and collections of the authors; queen (MNHN 1598).

Museum material

FRANCE: MHNG-FC: 47 specimens, Orange, 14 Sep. 1883. – MCSN-EC: Orange, label with bad handwriting which could be by Forel. – MNHN-BAC: Mont Ventoux, S face, Drôme, 16 Sep. 1978, coll.

Casevitz-Weulersse; 8 workers (2 pins), Gréolières, Alpes-Maritimes; ~30 workers (9 pins), Plateau de Caussols. – MCZL-KC: 5 pins, Caussol, S France, 9 Jun. 1954, leg. Stumper.

New material

FRANCE: All in personal collections of CG and RB: Plateau de Calern, 22 workers, 43.7553° N, 6.9113° E, leg. C. Galkowski, Aug. 2012, leg. A. Touchard; Plateau de Caussols, 23 workers, 43.7319° N, 6.9426° E, leg. C. Galkowski, Jun. 2012, leg. A. Touchard; Gréolières, 14 workers, 43.82° N, 6.92° E, Aug. 2012, leg. A. Touchard; Orange, 13 workers, 2 queens, 44.1344° N, 4.8084° E, 14 Jul. 2011, leg. R. Blatrix; Sisteron, 12 workers, 44.1989° N, 5.9419° E, 17 Jun. 2012, leg. R. Blatrix; Mont Ventoux, 41 workers, 4 queens, 44.1519° N, 5.3226° E, 2009, leg. C. Galkowski, 14 Jul. 2011, leg. R. Blatrix, Aug. 2011, leg. C. Lebas.

Description

Minor worker (n = 72)

CW 540–900 μm ; nCH 0–10 (4.5); nCU 0–8 (3.4); nTx 3–23 (12.9); nSc 0–7 (3.4); nG 1–27 (10.1); PDG 34–48 (40.3); GHL 70–170 μm (123.7); GHL/CW 0.124–0.263 (0.171).

Media and major worker (n = 53)

CW 900–1530 μm ; nCH 2–15 (8.4); nCU 2–17 (7.1); nTx 20–75 (38.9); nSc 3–22 (7.6); nG 12–43 (24.8); PDG 24–68 (43.7); GHL 140–240 μm (178.9); GHL/CW 0.117–0.186 (0.145).

Body black; only tibiae, scape and mandible brown. All parts of body with long erect hairs (GHL/CW > 0.12). Pubescence on dorsal surface of first gaster tergite sparse in all worker categories (PDG > 24), revealing smooth and shining cuticle. Profile of mesosoma sinuous. Petiolar scale erect, thick, slightly notched at summit in large workers. Head of minor workers clearly elongate, but less than in *P. nasuta* (CL/CW 1.16–1.28). Head of media and major workers even less elongate (CL/CW 1.046–1.19). Clypeus finely striate longitudinally, with faint trace of median carina. Frontal triangle and space between frontal carina also finely striate. Sculpture disappears toward occiput, cuticle becoming smooth and shining, or faintly punctuated in large workers. Scape long, surpassing occipital border.

Queen (n = 6)

CW 1530–1770 μm (1690); CL 1550–1750 μm (1660); SL 1220–1390 μm (1300); nCH 4–13 (8.20); nCU 5–7 (6.20); nTX 62–92 (79); nSc 8–13 (11.2); nG 26–36 (31.5); PDG 12–17 (14.5); GHL 210–240 μm (227); MW 1280–1410 μm (1330); ML 2240–2510 μm (2390); ScW 760–850 μm (820); GHL/CW 0.124–0.141 (0.135).

Color as in worker. Many long and erect hairs on all body parts (nG > 26, GHL > 200 μm). Some erect hairs also on femora and tibiae. Dense pubescence on entire body, masking surface of cuticle. Mesosoma less wide than head. Wing remains indicate winged queens, although wings possibly small given reduced development of scutum and scutellum. Petiolar scale high and wide, deeply notched at summit. Gaster rather small. Head almost as large as long, entirely and finely sculptured, faint riddles of anterior part replaced by puncture on posterior part, giving head an almost dull aspect. Clypeus finely striate longitudinally. Scape surpassing occipital border.

Male

Unknown.

Distribution

Southern France (Fig. 1).




Fig. 9. *Proformica longipilosa* sp. nov., holotype, minor worker from colony Mont Ventoux 10, France. **A.** Lateral view. **B.** Dorsal view. **C.** Head in full face view. **D.** Gaster in dorsal view. Scale bars = 1 mm. Automontage: Claude Lebas.

Remarks

We made direct comparisons between specimens of *P. longipilosa* sp. nov. and *P. longiseta* (A. Tinaut leg.) from Sierra Nevada, Spain, and *P. ferreri* (IRSNB-BC, 2 workers, 1 ♂ (type specimen) from Spain). The latter two species, in addition to *P. nasuta*, were formally described from western Europe. Specimens of *P. longipilosa* sp. nov. are unambiguously distinguished from those of these two species by the combination of the following characters: erect hairs on the body are longer ($GHL/CW > 0.12$) and the pubescence on the dorsal surface of the first gaster tergite is sparser. This last character is especially discriminant in media and major workers (PDG 24–68, mean = 43.7). In addition, the cuticle is smoother, giving a shinier appearance, in particular on the head.

Discussion

Based on arguments from morphometric analysis, DNA sequences and cuticular hydrocarbons, we show that the populations of *Proformica* in southern France belong to two species, one of which is *P. nasuta*, and the other a new species that we name *P. longipilosa* sp. nov.



Fig. 10. *Proformica longipilosa* sp. nov., queen from colony Mont Ventoux 10, France. **A.** Lateral view. **B.** Dorsal view. **C.** Head in full face view. Scale bars = 1 mm. Automontage: Claude Lebas.

Consequences for the taxonomy of *Proformica*

There has been much confusion in the concept of the taxon *P. nasuta*, in part because the type appears to have been lost a long time ago. None of the taxonomic studies published after the description of the species by Nylander (1856) made reference to the type, which was collected in Beaucaire (France). Forel (1886) described the worker and the queen from specimens collected in Orange (France) and sent samples to many of his colleagues throughout Europe and Russia. We examined the specimens from Orange in the Forel, Emery and Bondroit collections, and we collected new samples from the same locality in 2011. All differ markedly from those of Beaucaire and belong to the new species we describe in this study, *P. longipilosa* sp. nov. All the taxonomic studies after 1886 used the samples from Orange, or descriptions of them, as a reference for *P. nasuta*. These studies described *P. nasuta* as having long erect hairs and sparse pubescence (e.g., Ruzsky 1905; Emery 1909; Wheeler 1913; Bondroit 1918; Santschi 1925; Bernard 1968; Dlussky 1969; Collingwood 1976; Agosti & Collingwood 1987), two characters that are typical of *P. longipilosa* sp. nov. Other specimens from southeastern France (Plateau de Caussols, Tourettes-sur-Loup) have also been used as references for *P. nasuta* (Collingwood 1956; Stumper 1957; Dlussky 1969), but they come from an area that we now recognise as belonging to the range of *P. longipilosa* sp. nov., and are thus likely distinct from *P. nasuta*. This mistake has been perpetuated so that the actual conception of *P. nasuta* refers to *P. longipilosa* sp. nov. A consequence of this is that all reports of *P. nasuta* since 1886, including all those from eastern Europe and Asia, are probably erroneous.

There is a striking similarity between the biogeographic patterns of the genus *Proformica* and those observed for other steppe elements (Ruano *et al.* 2011; Sanllorente *et al.* 2015) such as, for instance, the meadow and steppe vipers (*Vipera ursinii* species complex). Both *Proformica* and the *V. ursinii* complex are known to occupy the same type of habitats and show the same pattern of disjunct distribution, with one area across Asia and one area restricted to western Europe. Moreover, they are both poor dispersers (Sanllorente *et al.* 2015; Ferchaud *et al.* 2011). It is likely that the current distribution and speciation patterns of the two groups have been induced by the same climatic and geographic events. The eastern and western clades of the *Vipera ursinii* species complex diverged about 4 Mya (Zinenko *et al.* 2015), and the taxa from the two geographic areas are completely distinct. In addition, mutation rates and the number of generations per unit time are expected to be higher in insects than in snakes (Martin & Palumbi 1993). We thus expect that the *Proformica* species from western Europe are distinct from those from the East.

Intraspecific variation

Localities for *P. nasuta* can be divided into two subgroups. One is composed of lowland localities that can be close to each other and form an almost continuous distribution in the plain of the Languedoc and the Rhône valley (Montpellier, Sauteyrargues, Pompignan, Grospierres, Collias, Beaucaire, Jonquières, Tarascon, Plaine de la Crau, Aurons and Bonnieux). The other subgroup is composed of localities isolated on the summits of medium-sized mountains (Sumène, Sainte-Baume, Grand Luberon and Montagne de Lure). Beyond ecological differences (mountain vs lowland), the distinction between the two groups is supported by the analysis of morphological characters, DNA and cuticular hydrocarbons. Except for Sumène, individuals from mountain localities have some erect hairs on the mesosoma, whereas hairs are lacking in most lowland individuals. Mountain individuals also have longer hairs and sparser pubescence on the gaster compared to lowland individuals. Interestingly, mitochondrial sequences from Sainte-Baume, Grand Luberon and Montagne de Lure form three monophyletic clades that are highly differentiated from each other, whereas most of the lowland localities show little differentiation. This pattern is consistent with a particularly high degree of isolation of the mountain localities, as already shown for *P. longiseta* in Spain (Sanllorente *et al.* 2015). Cuticular hydrocarbons also differ between lowland and mountain localities. At this stage we are reluctant to consider the mountain localities as

a separate species because lowland and mountain localities form a consistent group morphologically and do not form two monophyletic clades in the mitochondrial phylogenies. Instead, we consider that the population of each mountain locality is derived independently from the lowland population. Sanlloriente *et al.* (2015) proposed that climate-driven range fluctuation of populations of *P. longiseta* during Pleistocene glaciations induced a strong isolation among populations that are now restricted to mountain tops in southern Spain, because this species is adapted to cold and arid environments. Extant populations would be derived from an ancient, large population, independently from each other. A similar process may explain the divergence we noted between lowland and mountain populations of *P. nasuta*, and among mountain populations, and may have induced local adaptation of mountain populations. The morphological features of the mountain individuals of *P. nasuta* make them more similar to *P. longipilosa* sp. nov., especially where localities of the two species are close to each other (Grand Luberon and Montagne de Lure). These features might be the result either of introgression of *P. longipilosa* sp. nov. into populations of *P. nasuta*, or of morphological convergence in response to a similar environment.

Proformica longipilosa sp. nov. also shows two subgroups. One is composed of populations from localities on plateaus and mountains (Plateau de Calern, Plateau de Caussols, Gréolières and Mont Ventoux) and the other is composed of lower elevation localities (Orange, Sisteron and Vinsobres). Individuals of the lowland localities tend to have shorter hairs and denser pubescence on the gaster than those from high-elevation localities, and thus are morphologically closer to *P. nasuta* than the high-elevation individuals. As proposed for *P. nasuta*, these features could result from either introgression or convergence. Although lowland *P. longipilosa* sp. nov. and mountain *P. nasuta* tend to converge morphologically, they can still be easily distinguished, leaving no doubt regarding their assignment to species. All nests from the three low-elevation localities of *P. longipilosa* sp. nov. had mitochondrial sequences typical of *P. nasuta*. We hypothesise that this incongruence between morphological and molecular characters, specifically for low-elevation localities, results from introgression of mitochondrial DNA from *P. nasuta* to *P. longipilosa* sp. nov. Such introgression would be most likely to occur in low-elevation localities because *P. nasuta* is relatively widespread in the lowlands. Complete or gene-specific introgression of maternal DNA is a well-known phenomenon in insects (Ballard 2000; Chan & Levin 2005; Linnen & Farrell 2007). An isolated event of partial mitochondrial introgression is also suggested in Plateau de Calern, where all nests were morphologically classified as *P. longipilosa* sp. nov., but one had *Cytb* sequences typical of *P. nasuta*. All other nests in this locality fitted within *P. longipilosa* sp. nov. for both mitochondrial markers. It is worth noting that we could not find any locality with nests of both *P. nasuta* and *P. longipilosa* sp. nov., although the two can be found in similar habitats.

Interpretation of cuticular hydrocarbons

Cuticular hydrocarbons separate the two species without ambiguity. Populations from all localities of *P. nasuta* form a homogeneous clade with relatively little differentiation except for two localities isolated at the summits of mountains (Grand Luberon and Montagne de Lure). *Proformica longipilosa* sp. nov. (Mont Ventoux and Sisteron) appears to be well separated from all other ants studied here, including the outgroups (*P. longiseta* and *Cataglyphis* Förster, 1850), confirming its status as a separate species. This classification is globally consistent with spatial distribution of the localities and with the classification based on morphology and DNA sequences. Localities of *P. nasuta* from the lowlands form continuous populations without important geographical isolation, allowing regular exchange of migrants resulting in little differentiation of cuticular hydrocarbons. However, the mountain localities Grand Luberon and Montagne de Lure are isolated and, probably as a consequence, are divergent for cuticular hydrocarbons. On the mitochondrial tree they also diverge from other localities of *P. nasuta*. Cuticular hydrocarbons in *Proformica* appear to be linked to phylogenetic signature but seem to change rapidly with geographical isolation, even faster than mitochondrial DNA. Geographic variation in cuticular hydrocarbons depends on the taxon. For instance, profiles are very stable for *Formica* ants from Finland to Great Britain (Martin

et al. 2008) and for *Lasius niger* Linnaeus, 1758 from Denmark to the south of France (Lenoir *et al.* 2009; Lenoir unpubl.). In contrast, rapid spatial changes in hydrocarbons are present in some taxa like *Odontomachus* Latreille, 1804 (Smith *et al.* 2013) and *Cataglyphis* (Dahbi *et al.* 1996). Interestingly, *Rossomyrmex minuchae* Tinaut, 1981, a slave-maker parasite of *Proformica longiseta*, also has different chemotypes in three populations in Sierra Nevada, Spain (Sanllorente *et al.* 2012). It is noteworthy that the genus *Cataglyphis* is phylogenetically, biologically and ecologically very close to *Proformica*. Both have very limited queen dispersal, are specialized on dry habitats and forage on dead invertebrates at the warmest time of the day. It would be worth investigating whether strong divergence in cuticular hydrocarbons within species could be related to one or more of these characteristics.

Conclusions

Although we relied on an integrative taxonomy approach, using several complementary sources of information, we confirm the general view that the taxonomy of the genus *Proformica* is a complex problem. The nuclear markers chosen for use here evolve too slowly and thus lack resolution. Information from mitochondrial genes is blurred by genetic processes such as transposition and introgression and may be biased by queen philopatry. Our results suggest that morphology is a better tool to resolve taxonomy in this genus than either cuticular hydrocarbons or DNA sequences of the genes commonly used for phylogenetic analyses and barcoding, although genetic markers other than those used in this study should also be investigated. However, for the genus *Proformica*, the zoogeographical region of southern France is the least complex in taxonomic terms. Thus, the morphological approach developed in this paper may prove unreliable in other regions, such as the Iberian Peninsula and eastern Europe. A population-genetic approach using tools such as microsatellites or single nucleotide polymorphisms from Next Generation Sequencing (e.g., RADseq) may help disentangle this taxonomic knot.

Acknowledgements

This project was supported by the network “Bibliothèque du Vivant” funded by CNRS, MNHN, INRA and the CEA (Centre National de Séquençage). We thank Axel Touchard, Théotime Colin, Philippe Geniez and V. Bouchet for providing ant samples, Nicolas Lescureux for help with translation of papers in Russian, Hugo Darras, Marek Borowiec, Birgit Schlick-Steiner and Omid Paknia for useful comments on an earlier version of the manuscript, Alberto Tinaut and Xavier Espadaler for constructive remarks, Alberto Tinaut for sending specimens of *P. longiseta* from Sierra Nevada, Doyle McKey and Finn Kjellberg for English editing, Agnièle Touret-Alby and Quentin Rome (MNHN) for data on the specimens of *Proformica* in the Paris collections and for the loan of specimens, and Bernhard Merz (Entomology Department, MHNG), Wouter Dekoninck (Entomology Department, IRSNB), Isabelle Zürcher-Pfander (NMB), Roberto Poggi (MCSN) and Anne Freitag (MCZL) for communicating data regarding the collections of A. Forel, J. Bondroit, F. Santschi, C. Emery and H. Kutter, respectively.

References

- Agosti D. 1994. The phylogeny of the ant tribe Formicini (Hymenoptera, Formicidae) with the description of a new genus. *Systematic Entomology* 19: 93–117. <https://doi.org/10.1111/j.1365-3113.1994.tb00581.x>
- Agosti D. & Collingwood C.A. 1987. A provisional list of the Balkan ants (Hym. Formicidae) with a key to the worker caste. II. Key to the worker caste, including the European species without the Iberian. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 60: 261–293.
- Ballard J.W.O. 2000. When one is not enough: Introgression of mitochondrial DNA in *Drosophila*. *Molecular Biology and Evolution* 17: 1126–1130. <https://doi.org/10.1093/oxfordjournals.molbev.a026394>

- Belshaw R. & Quicke D.L.J. 1997. A molecular phylogeny of the Aphidiinae (Hymenoptera: Braconidae). *Molecular Phylogenetics and Evolution* 7: 281–293. <https://doi.org/10.1006/mpev.1996.0400>
- Bernard F. 1968. *Faune de l'Europe et du Bassin Méditerranéen. 3. Les Fourmis (Hymenoptera Formicidae) d'Europe occidentale et septentrionale*. Masson, Paris.
- Bolton B. 2014. An Online Catalog of the Ants of the World. Available from <http://antcat.org> [accessed 26 Jan. 2017].
- Bondroit J. 1918. Les fourmis de France et de Belgique. *Annales de la Société Entomologique de France* 87: 1–174.
- Chan K.M.A. & Levin S.A. 2005. Leaky prezygotic isolation and porous genomes: Rapid introgression of maternally inherited DNA. *Evolution* 59: 720–729. <https://doi.org/10.1554/04-534>
- Collingwood C.A. 1956. Ant hunting in France. *L'entomologiste* 89: 105–108.
- Collingwood C.A. 1976. A provisional list of Iberian Formicidae with a key to the worker caste. *EOS – Revista Española de Entomología* 52: 65–95.
- Collingwood C.A. & Yarrow I.H.H. 1969. A survey of Iberian Formicidae. *EOS – Revista Espanola de Entomologia* 44: 53–101.
- Dahbi A., Cerda X., Hefetz A. & Lenoir A. 1996. Social closure, aggressive behavior, and cuticular hydrocarbon profiles in the polydomous ant *Cataglyphis iberica* (Hymenoptera, Formicidae). *Journal of Chemical Ecology* 22: 2173–2186. <https://doi.org/10.1007/BF02029538>
- Dlussky G.M. 1969. Ants of the genus *Proformica* Ruzs. of the USSR and contiguous countries (Hymenoptera: Formicidae). *Zoologicheskii Zhurnal* 48: 217–232. [in Russian]
- Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Emery C. 1909. Beiträge zur Monographie der Formiciden des paläarktischen Faunengebietes (Hym.). Teil VII. *Deutsche Entomologische Zeitschrift* 1909: 179–204.
- Emery C. 1925. Hymenoptera, Fam. Formicidae, subfam. Formicinae. *Genera Insectorum* 183: 1–302.
- Errard C., Ruano F., Richard F.J., Lenoir A., Tinaut A. & Hefetz A. 2006. Co-evolution-driven cuticular hydrocarbon variation between the slave-making ant *Rossomyrmex minuchae* and its host *Proformica longiseta* (Hymenoptera: Formicidae). *Chemoecology* 16: 235–240. <https://doi.org/10.1007/s00049-006-0358-4>
- Espadaler X. & Cagniant H. 1987. Contribution à la connaissance des fourmis marocaines. Description du mâle de *Proformica theryi* Santschi, 1936 (Hymenoptera, Formicidae). *Nouvelle Revue d'Entomologie* 4: 133–138.
- Ferchaud A.L., Lyet A., Cheylan M., Arnal V., Baron J.P., Montgelard C. & Ursenbacher S. 2011. High genetic differentiation among French populations of the Orsini's Viper (*Vipera ursinii ursinii*) based on mitochondrial and microsatellite data: implications for conservation management. *Journal of Heredity* 102: 67–78. <https://doi.org/10.1093/jhered/esq098>
- Forel A. 1886. Études myrmécologiques en 1886. *Annales de la Société Entomologique de Belgique* 30: 131–215.
- Goropashnaya A.V., Fedorov V.B., Seifert B. & Pamilo P. 2004. Limited phylogeographical structure across Eurasia in two red wood ant species *Formica pratensis* and *F. lugubris* (Hymenoptera, Formicidae). *Molecular Ecology* 13: 1849–1858. <https://doi.org/10.1111/j.1365-294X.2004.02189.x>

- Goropashnaya A.V., Fedorov V.B., Seifert B. & Pamilo P. 2007. Phylogeography and population structure in the ant *Formica exsecta* (Hymenoptera, Formicidae) across Eurasia as reflected by mitochondrial DNA variation and microsatellites. *Annales Zoologici Fennici* 44: 462–474.
- Goropashnaya A.V., Fedorov V.B., Seifert B. & Pamilo P. 2012. Phylogenetic relationships of Palaearctic *Formica* species (Hymenoptera, Formicidae) based on mitochondrial cytochrome *b* sequences. *PLoS ONE* 7: e41697. <https://doi.org/10.1371/journal.pone.0041697>
- Hasegawa E., Tinaut A. & Ruano F. 2002. Molecular phylogeny of two slave-making ants: *Rossomyrmex* and *Polyergus* (Hymenoptera: Formicidae). *Annales Zoologici Fennici* 39: 267–271.
- Hebert P.D.N., Penton E.H., Burns J.M., Janzen D.H. & Hallwachs W. 2004. Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences* 101: 14812–14817. <https://doi.org/10.1073/pnas.0406166101>
- Lanfear R., Calcott B., Ho S.Y.W. & Guindon S. 2012. PartitionFinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1695–1701. <https://doi.org/10.1093/molbev/mss020>
- Lenoir A., Depickere S., Devers S., Christides J.P. & Detrain C. 2009. Hydrocarbons in the ant *Lasius niger*: from the cuticle to the nest and home range marking. *Journal of Chemical Ecology* 35: 913–921. <https://doi.org/10.1007/s10886-009-9669-6>
- Liautard C. & Keller L. 2001. Restricted effective queen dispersal at a microgeographic scale in polygynous populations of the ant *Formica exsecta*. *Evolution* 55: 2484–2492. <https://doi.org/10.1111/j.0014-3820.2001.tb00763.x>
- Linnen C.R. & Farrell B.D. 2007. Mitonuclear discordance is caused by rampant mitochondrial introgression in *Neodiprion* (Hymenoptera: Diprionidae) sawflies. *Evolution* 61: 1417–1438. <https://doi.org/10.1111/j.1558-5646.2007.00114.x>
- Martin A.P. & Palumbi S.R. 1993. Body size, metabolic rate, generation time, and the molecular clock. *Proceedings of the National Academy of Sciences* 90: 4087–4091. <https://doi.org/10.1073/pnas.90.9.4087>
- Martin S.J., Helanterä H. & Drijfhout F.P. 2008. Evolution of species-specific cuticular hydrocarbon patterns in *Formica* ants. *Biological Journal of the Linnean Society* 95: 131–140. <https://doi.org/10.1111/j.1095-8312.2008.01038.x>
- Moreau C.S., Bell C.D., Vila R., Archibald S.B. & Pierce N.E. 2006. Phylogeny of the ants: diversification in the age of angiosperms. *Science* 312: 101–104. <https://doi.org/10.1126/science.1124891>
- Nowbahari E., Lenoir A., Clément J.L., Lange C., Bagnères A.G. & Joulie C. 1990. Individual, geographical and experimental variation of cuticular hydrocarbons of the ant *Cataglyphis cursor* (Hymenoptera: Formicidae): their use in nest and subspecies recognition. *Biochemical Systematics and Ecology* 18: 63–73. [https://doi.org/10.1016/0305-1978\(90\)90036-F](https://doi.org/10.1016/0305-1978(90)90036-F)
- Nylander W. 1856. Synopsis des formicides de France et d'Algérie. *Annales des Sciences Naturelles: Zoologie et Biologie Animale*, 4th Series 5: 51–109. Available from <http://biodiversitylibrary.org/page/13469211> [accessed 12 Feb. 2017].
- Oppelt A., Spitzenpfeil N., Kroiss J. & Heinze J. 2008. The significance of intercolonial variation of cuticular hydrocarbons for inbreeding avoidance in ant sexuals. *Animal Behaviour* 76: 1029–1034. <https://doi.org/10.1016/j.anbehav.2008.05.020>
- Radchenko A. 2007. The ants (Hymenoptera, Formicidae) in the collection of William Nylander. *Fragmenta Faunistica* (Warsaw) 50: 27–41.
- Reyment R.A. 1989. Compositional data analysis. *Terra Review* 1: 29–34.

- Ronquist F., Teslenko M., van der Mark P., Ayres D.L., Darling A., Höhna S., Larget B., Liu L., Suchard M.A. & Huelsenbeck J.P. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Ruano F., Devers S., Sanllorrente O., Errard C., Tinaut A. & Lenoir A. 2011. A geographical mosaic of coevolution in a slave-making host–parasite system. *Journal of Evolutionary Biology* 24: 1071–1079. <https://doi.org/10.1111/j.1420-9101.2011.02238.x>
- Ruzsky M. 1905. The ants of Russia. *Trudy Obshchestva Estestvoispytatelei pri Imperatorskom Kazanskom Universitete* 38: 1–800. [in Russian]
- Sanllorrente O., Lorite P., Devers S., Ruano F., Lenoir A. & Tinaut A. 2012. The spatial distribution does not affect host–parasite coevolution in *Rossomyrmex* ants. *Insectes Sociaux* 59: 361–368. <https://doi.org/10.1007/s00040-012-0228-8>
- Sanllorrente O., Ruano F. & Tinaut A. 2015. Large-scale population genetics of the mountain ant *Proformica longiseta* (Hymenoptera: Formicidae). *Population Ecology* 57: 637–648. <https://doi.org/10.1007/s10144-015-0505-2>
- Santschi F. 1925. Fourmis d’Espagne et autres espèces paléarctiques (Hymenopt.). *EOS – Revista Española de Entomología* 1: 339–360.
- Saux C., Fisher B.L. & Spicer G.S. 2004. Dracula ant phylogeny as inferred by nuclear 28S rDNA sequences and implications for ant systematics (Hymenoptera: Formicidae: Amblyoponinae). *Molecular Phylogenetics and Evolution* 33: 457–468. <https://doi.org/10.1016/j.ympev.2004.06.017>
- Schlick-Steiner B.C., Steiner F.M., Seifert B., Stauffer C., Christian E. & Crozier R.H. 2010. Integrative taxonomy: a multisource approach to exploring biodiversity. *Annual Review of Entomology* 55: 421–438. <https://doi.org/10.1146/annurev-ento-112408-085432>
- Seifert B. 2007. *Die Ameisen Mittel- und Nordeuropas*. Lutra, Görlitz, Germany.
- Smith A.A., Millar J.G., Hanks L.M. & Suarez A.V. 2013. A conserved fertility signal despite population variation in the cuticular chemical profile of the trap-jaw ant *Odontomachus brunneus*. *Journal of Experimental Biology* 216: 3917–3924. <https://doi.org/10.1242/jeb.089482>
- Stamatakis A., Hoover P. & Rougemont J. 2008. A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology* 57: 758–771. <https://doi.org/10.1080/10635150802429642>
- Stumper R. 1957. Sur l’éthologie de la fourmi a miel *Proformica nasuta* Nyl. (Étude myrmécologique LXXVIII). *Bulletin de la Société des Naturalistes Luxembourgeois* 60: 87–97.
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M. & Kumar S. 2011. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Ward P.S. & Downie D.A. 2005. The ant subfamily Pseudomyrmecinae (Hymenoptera: Formicidae): phylogeny and evolution of big-eyed arboreal ants. *Systematic Entomology* 30: 310–335. <https://doi.org/10.1111/j.1365-3113.2004.00281.x>
- Wheeler W.M. 1913. A revision of the ants of the genus *Formica* (Linné) Mayr. *Bulletin of the Museum of Comparative Zoology* 53: 379–565.
- Zinenko O., Stümpel N., Mazanaeva L., Bakiev A., Shiryaev K., Pavlov A., Kotenko T., Kukushkin O., Chikin Y., Duisebajeva T., Nilson G., Orlov N.L., Tuniyev S., Ananjeva N.B., Murphy R.W. & Joger U. 2015. Mitochondrial phylogeny shows multiple independent ecological transitions and northern

dispersion despite of Pleistocene glaciations in meadow and steppe vipers (*Vipera ursinii* and *Vipera renardi*). *Molecular Phylogenetics and Evolution* 84: 85–100. <https://doi.org/10.1016/j.ympev.2014.12.005>

Manuscript received: 1 March 2016

Manuscript accepted: 31 January 2017

Published on: 2 March 2017

Guest editors: Line Le Gall, Frédéric Delsuc, Stéphane Hourdez, Guillaume Lecointre and Jean-Yves Rasplus

Desk editor: Danny Eibye-Jacobsen

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands.

Appendices

Appendix 1. List of the nest samples collected for this study, with geographic locality, number of individuals per nest used for morphological analyses and GenBank accession numbers for DNA markers.

colony#	species	locality	municipality	country	date	latitude	longitude	altitude_m	collector	morpho_workers	morpho_queens	COI
Cytb 28S LwRh												
RB0000530	<i>Cataglyphis cursor</i>	Riboux (83)	France	3Jun.2011	43.32693	5.76608	Rumsais Blatrix	KU749638	KU749738			
RB0000577	<i>Formica cunicularia</i>	Estoher (66)	France	25Jun.2011	42.574	2.48461	Rumsais Blatrix	KU749784	KU749819			
RB0000971	<i>Cataglyphis mauritanica</i>	Morocco		27May2012	33.3818	-5.1324	Philippe Geniez	KU749639	KU749785	KU749820		
RB0001590	<i>Bajcaridris theryi</i>	Morocco		Jul.2012	Christophe Galkowski		KU749786	KU749821				
Salon 1	<i>Cataglyphis cursor</i>	Salon-de-Provence (13)	France	Jul.2012	43.63705	5.02083	Claude Lebas	KU749640	KU749787	KU749822		
Aurons 1	<i>Proformica nasuta</i>	Aurons Aurons (13)	France	May2006	43.66	5.15	199 V. Bouchet	4				
Beaucaire 1	<i>Proformica nasuta</i>	Beaucaire Beaucaire (30)	France	9Jul.2011	43.83544	4.61828	21 Rumsais Blatrix, Claude Lebas	4	2			
KU749600												
Beaucaire 2	<i>Proformica nasuta</i>	Beaucaire Beaucaire (30)	France	9Jul.2011	43.83547	4.61858	19 Rumsais Blatrix, Claude Lebas	3				
KU749656												
Beaucaire 3	<i>Proformica nasuta</i>	Beaucaire Beaucaire (30)	France	9Jul.2011	43.83533	4.61878	17 Rumsais Blatrix, Claude Lebas	3				
KU749601, KU749602												
Beaucaire 4	<i>Proformica nasuta</i>	Beaucaire Beaucaire (30)	France	9Jul.2011	43.83531	4.61928	15 Rumsais Blatrix, Claude Lebas	4				
KU749603, KU749604, KU749605												
Bonnieux 1	<i>Proformica nasuta</i>	Bonnieux Bonnieux (84)	France	1Oct.2011	43.8625	5.30694	166 Claude Lebas	4	KU749606	KU749659		
KU749759												
Bonnieux 2	<i>Proformica nasuta</i>	Bonnieux Bonnieux (84)	France	1Oct.2011	43.8625	5.30694	166 Claude Lebas	4	KU749660			
Plateau de Calern 10	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	19Aug.2010	43.7514	6.92349	1264 Christophe Galkowski	4				
Plateau de Calern 18	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	19Aug.2010	43.7514	6.92349	1264 Christophe Galkowski	4				
Plateau de Calern 2	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	Aug.2012	43.75325	6.9159	1256 Axel Touchard	3	KU749607	KU749661	KU749760	KU749795
Plateau de Calern 3	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	Aug.2012	43.75457	6.91025	1272 Axel Touchard	3	KU749608	KU749662	KU749761	KU749796
Plateau de Calern 4	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Cipières (06)	France	Aug.2012	43.75678	6.90642	1283 Axel Touchard	3	KU749609	KU749663	KU749762	KU749797
Plateau de Calern 6	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Andon (06)	France	Aug.2012	43.77595	6.85128	1127 Axel Touchard	3	KU749610	KU749664	KU749763	KU749798
Plateau de Calern 7	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	19.Aug2010	43.7514	6.92349	1264 Christophe Galkowski	2	KU749611, KU749612	KU749665, KU749666	KU749764	KU749799
Plateau de Calern 8	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	19.Aug2010	43.7514	6.92349	1264 Christophe Galkowski	KU749667				
Plateau de Calern 9	<i>Proformica longipilosa</i> sp. nov.	Plateau de Calern Caussols (06)	France	19.Aug2010	43.7514	6.92349	1264 Christophe Galkowski	KU749668				
Plateau de Caussols 1	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	Jun.2012	43.73504	6.94663	1102 Axel Touchard	3	KU749672			
Plateau de Caussols 10	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	19Aug.2010	43.7331	6.94244	1115 Christophe Galkowski	KU749669				
Plateau de Caussols 11	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	19Aug.2010	43.7331	6.94244	1115 Christophe Galkowski	KU749613	KU749670	KU749765	KU749800	
Plateau de Caussols 19	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	19Aug.2010	43.7331	6.94244	1115 Christophe Galkowski	3	KU749671			
Plateau de Caussols 2	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	Jun.2012	43.73497	6.94676	1100 Axel Touchard	3	KU749673			
Plateau de Caussols 3	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	Jun.2012	43.73488	6.94681	1100 Axel Touchard	3	KU749674			
Plateau de Caussols 4	<i>Proformica longipilosa</i> sp. nov.	Plateau de Caussols Gourdon (06)	France	Jun.2012	43.73493	6.94699	1098 Axel Touchard	2				

- Plateau de Caussols 5 *Proformica longipilosa* sp. nov. Plateau de Caussols Caussols (06) France Aug.2012 43.7219 6.90923 1124 Axel Touchard 3 KU749614 KU749675 KU749766 KU749801
- Plateau de Caussols 6 *Proformica longipilosa* sp. nov. Plateau de Caussols Caussols (06) France Aug.2012 43.72737 6.93098 1139 Axel Touchard 3 KU749615 KU749676 KU749767 KU749802
- Plateau de Caussols 7 *Proformica longipilosa* sp. nov. Plateau de Caussols Gourdon (06) France Aug.2012 43.73065 6.97157 1033 Axel Touchard 3 KU749677
- Collias 1 *Proformica nasuta* Collias Collias (30) France 12Jun.2010 43.947721 4.462309 162 Rumsais Blatrix 5
- Plaine de la Crau 1 *Proformica nasuta* Plaine de la Crau Saint-Martin-de-Crau (13) France Jun.2011 43.58333 4.83333 19 Claude Lebas 5 KU749616 KU749678 KU749768 KU749803
- Plaine de la Crau 2 *Proformica nasuta* Plaine de la Crau Saint-Martin-de-Crau (13) France Aug.2011 43.58333 4.83333 19 Claude Lebas 5
- Plaine de la Crau 3 *Proformica nasuta* Plaine de la Crau Saint-Martin-de-Crau (13) France Aug.2011 43.58333 4.83333 19 Claude Lebas 5 KU749679
- Plaine de la Crau 4 *Proformica nasuta* Plaine de la Crau Saint-Martin-de-Crau (13) France Aug.2011 43.58333 4.83333 19 Claude Lebas 3
- Gréolières 7 *Proformica longipilosa* sp. nov. Gréolières Gréolières (06) France Aug.2012 43.82862 6.94738 1409 Axel Touchard 3 KU749617 KU749680 KU749769 KU749804
- Gréolières 8 *Proformica longipilosa* sp. nov. Gréolières Gréolières (06) France Aug.2012 43.81688 6.8898 1202 Axel Touchard 3 KU749770 KU749805
- Grospierres 1 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749682
- Grospierres 10 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749681
- Grospierres 2 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749618 KU749683 KU749771 KU749806
- Grospierres 3 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749684
- Grospierres 4 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749619 KU749685 KU749772 KU749807
- Grospierres 5 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749686
- Grospierres 6 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749687
- Grospierres 7 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749688
- Grospierres 8 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 Théotime Colin 3 KU749689
- Grospierres 9 *Proformica nasuta* Grospierres Grospierres (07) France 6Jul.2012 44.41163 4.27133 102 ThÃ©otime Colin 3 KU749690
- Jonquières 1 *Proformica nasuta* Jonquières Jonquières-Saint-Vincent (30) France 11May2011 43.83144 4.57653 21 Rumsais Blatrix 2 KU749620 KU749691, KU749692
- Jonquières 2 *Proformica nasuta* Jonquières Jonquières-Saint-Vincent (30) France 11May2011 43.83144 4.57653 21 Rumsais Blatrix 2 KU749621 KU749693 KU749773 KU749808
- Jonquières 3 *Proformica nasuta* Jonquières Jonquières-Saint-Vincent (30) France 11May2011 43.83144 4.57653 21 Rumsais Blatrix 3 KU749622 KU749694, KU749695
- Jonquières F *Proformica nasuta* Jonquières Jonquières-Saint-Vincent (30) France 11May2011 43.83144 4.57653 21 Rumsais Blatrix KU749696
- Grand Luberon 1 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81713 5.4782 1038 Rumsais Blatrix 3 KU749623 KU749697 KU749774 KU749809
- Grand Luberon 2 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81716 5.47816 1038 Rumsais Blatrix 4 KU749698
- Grand Luberon 3 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81689 5.47868 1040 Rumsais Blatrix 3 KU749699
- Grand Luberon 4 *Proformica nasuta* Grand Luberon Cabrières-d'Aigues (84) France 2Jun.2012 43.81662 5.478 1050 Rumsais Blatrix 3 KU749700
- Grand Luberon 5 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81677 5.47548 1091 Rumsais Blatrix 3 KU749701
- Grand Luberon 6 *Proformica nasuta* Grand Luberon Cabrières-d'Aigues (84) France 2Jun.2012 43.8155 5.47376 1085 Rumsais Blatrix 3 KU749702
- Grand Luberon 7 *Proformica nasuta* Grand Luberon Cabrières-d'Aigues (84) France 2Jun.2012 43.81543 5.47168 1086 Rumsais Blatrix 3 KU749703
- Grand Luberon 8 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81574 5.46942 1091 Rumsais Blatrix 3 KU749704
- Grand Luberon 9 *Proformica nasuta* Grand Luberon Castellet (84) France 2Jun.2012 43.81941 5.46737 1047 Rumsais Blatrix 3 KU749624 KU749705 KU749775 KU749810
- Montagne de Lure 1 *Proformica nasuta* Montagne de Lure Châteauneuf-Val-Saint-Donat (04) France 16Jun.2012 44.11319 5.88414 1416 Rumsais Blatrix 3 KU749706 KU749776 KU749811
- Montagne de Lure 2 *Proformica nasuta* Montagne de Lure Châteauneuf-Val-Saint-Donat (04) France 16Jun.2012 44.11317 5.88399 1417 Rumsais Blatrix 3 KU749707
- Montagne de Lure 3 *Proformica nasuta* Montagne de Lure Châteauneuf-Val-Saint-Donat (04) France 16Jun.2012 44.11314 5.88378 1419 Rumsais Blatrix 3 KU749708

- Montagne de Lure 4 *Proformica nasuta* Montagne de Lure Châteauneuf-Val-Saint-Donat (04) France 16Jun.2012 44.11312 5.8837 1419 Rumsais Blatrix 3 KU749709
- Montagne de Lure 5 *Proformica nasuta* Montagne de Lure Châteauneuf-Val-Saint-Donat (04) France 16Jun.2012 44.1131 5.88352 1419 Rumsais Blatrix 3 KU749710 KU749777 KU749812
- Montpellier 1 *Proformica nasuta* Montpellier Montpellier (34) France 8Mar.2012 43.62914 3.89072 38 Rumsais Blatrix 2 KU749711
- Montpellier 2 *Proformica nasuta* Montpellier Montpellier (34) France 8Mar.2012 43.62925 3.89072 40 Rumsais Blatrix 1 KU749712
- Orange 1 *Proformica longipilosa* sp. nov. Orange Orange (84) France 14Jul.2011 44.13486 4.80817 93 Rumsais Blatrix 8 2 KU749625, KU749626 KU749713 KU749778 KU749813
- Orange 2 *Proformica longipilosa* sp. nov. Orange Orange (84) France 14Jul.2011 44.1345 4.8084 96 Rumsais Blatrix 1
- Orange 3 *Proformica longipilosa* sp. nov. Orange Orange (84) France 14Jul.2011 44.13394 4.80864 99 Rumsais Blatrix 4 KU749627 KU749714 KU749779 KU749814
- Sainte-Baume 1 *Proformica nasuta* Sainte-Baume Riboux (83) France 3Jun.2011 43.32738 5.77002 950 Rumsais Blatrix 3 KU749628, KU749629, KU749630 KU749721, KU749722 KU749780 KU749815
- Sainte-Baume 11 *Proformica nasuta* Sainte-Baume Cuges-les-pins (13) France Jul.2012 43.31528 5.71057 1012 Claude Lebas 4 1 KU749715
- Sainte-Baume 12 *Proformica nasuta* Sainte-Baume Cuges-les-pins (13) France Jul.2012 43.31495 5.70795 1009 Claude Lebas 4 KU749716
- Sainte-Baume 13 *Proformica nasuta* Sainte-Baume Cuges-les-pins (13) France Jul.2012 43.31528 5.71057 1012 Claude Lebas 3 KU749717
- Sainte-Baume 14 *Proformica nasuta* Sainte-Baume Plan-d'Aups-Sainte-Baume (83) France Jul.2012 43.31582 5.70975 1017 Claude Lebas KU749718
- Sainte-Baume 15 *Proformica nasuta* Sainte-Baume Cuges-les-pins (13) France Jul.2012 43.31468 5.70517 996 Claude Lebas KU749719
- Sainte-Baume 16 *Proformica nasuta* Sainte-Baume Cuges-les-pins (13) France Jul.2012 43.31468 5.70517 996 Claude Lebas KU749720
- Sainte-Baume 2 *Proformica nasuta* Sainte-Baume Riboux (83) France 3Jun.2011 43.32738 5.77002 950 Rumsais Blatrix KU749632, KU749633 KU749725
- Sainte-Baume 21 *Proformica nasuta* Sainte-Baume Plan-d'Aups-Sainte-Baume (83) France Jul.2012 43.31937 5.68035 864 Claude Lebas KU749631 KU749723 KU749781 KU749816
- Sainte-Baume 22 *Proformica nasuta* Sainte-Baume Plan-d'Aups-Sainte-Baume (83) France Jul.2012 43.31937 5.68035 864 Claude Lebas 3 KU749724
- Sainte-Baume 3 *Proformica nasuta* Sainte-Baume Riboux (83) France 3Jun.2011 43.32738 5.77002 950 Christophe Galkowski 4 KU749726
- Sainte-Baume 4 *Proformica nasuta* Sainte-Baume Riboux (83) France 3Jun.2011 43.32738 5.77002 950 Christophe Galkowski 5 KU749727
- Sainte-Baume 5 *Proformica nasuta* Sainte-Baume Riboux (83) France Aug.2011 43.32738 5.77002 950 Claude Lebas 4 KU749728
- Sainte-Baume 6 *Proformica nasuta* Sainte-Baume Riboux (83) France Aug.2011 43.32738 5.77002 950 Claude Lebas 4 KU749729, KU749730
- Sainte-Baume 7 *Proformica nasuta* Sainte-Baume Riboux (83) France 10Sep.2011 43.32738 5.77002 950 Rumsais Blatrix 4 KU749731
- Sainte-Baume 8 *Proformica nasuta* Sainte-Baume Riboux (83) France 10Sep.2011 43.32738 5.77002 950 Rumsais Blatrix 4 KU749732
- Sainte-Baume 9 *Proformica nasuta* Sainte-Baume Riboux (83) France 10Sep.2011 43.32738 5.77002 950 Rumsais Blatrix 4 KU749733
- Pompignan 1 *Proformica nasuta* Pompignan Pompignan (30) France May2010 43.89747 3.82489 199 Rumsais Blatrix 1 KU749636 KU749735 KU749782 KU749817
- Pompignan 10 *Proformica nasuta* Pompignan Pompignan (30) France 10Jul.2011 43.899 3.82611 196 Rumsais Blatrix, Philippe Wegnez 2 KU749634, KU749635 KU749734
- Pompignan 2 *Proformica nasuta* Pompignan Pompignan (30) France May2010 43.89747 3.82489 199 Rumsais Blatrix 1 KU749637 KU749736, KU749737 KU749783 KU749818
- Pompignan 3 *Proformica nasuta* Pompignan Pompignan (30) France May2010 43.89747 3.82489 199 Rumsais Blatrix 1
- Sauteyrargues 1 *Proformica nasuta* Sauteyrargues Sauteyrargues (34) France 28May2011 43.8275 3.91917 118 Rumsais Blatrix 2 KU749641 KU749739 KU749788 KU749823
- Sauteyrargues 2 *Proformica nasuta* Sauteyrargues Sauteyrargues (34) France 28May2011 43.8275 3.91917 118 Rumsais Blatrix 3 KU749740
- Sisteron 1 *Proformica longipilosa* sp. nov. Sisteron Sisteron (04) France 17Jun.2012 44.19895 5.94151 554 Rumsais Blatrix 3 KU749642 KU749741 KU749789 KU749824
- Sisteron 2 *Proformica longipilosa* sp. nov. Sisteron Sisteron (04) France 17Jun.2012 44.19872 5.94217 558 Rumsais Blatrix 3 KU749742
- Sisteron 3 *Proformica longipilosa* sp. nov. Sisteron Sisteron (04) France 17Jun.2012 44.19898 5.9419 558 Rumsais Blatrix 3 KU749743
- Sisteron 4 *Proformica longipilosa* sp. nov. Sisteron Sisteron (04) France 17Jun.2012 44.1991 5.94187 558 Rumsais Blatrix 3 KU749643 KU749744 KU749790 KU749825
- Sumène 1 *Proformica nasuta* Montagne de la Fage Sumène (30) France 9Apr.2012 43.99044 3.7714 929 Rumsais Blatrix 3 KU749644 KU749745
- Sumène 2 *Proformica nasuta* Montagne de la Fage Sumène (30) France 9Apr.2012 43.99044 3.7714 929 Rumsais Blatrix 3 KU749746
- Tarascon 1 *Proformica nasuta* Tarascon Tarascon (13) France 9Jul.2011 43.84189 4.73778 15 Rumsais Blatrix, Claude Lebas 4 1 KU749645 KU749747
- Tarascon 2 *Proformica nasuta* Tarascon Tarascon (13) France 9Jul.2011 43.84186 4.73744 16 Rumsais Blatrix, Claude Lebas 4 KU749748

Tarascon 3 *Proformica nasuta* Tarascon Tarascon (13) France 9Jul.2011 43.84247 4.7395 16 Rumsais Blatrix, Claude Lebas 4 KU749646 KU749749

Mont Ventoux 1 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 2009 44.15158 5.32319 1466 Christophe Galkowski 3 KU749652 KU749756 KU749791 KU749826

Mont Ventoux 10 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 14Jul.2011 44.15261 5.32081 1442 Rumsais Blatrix 4 KU749647, KU749648 KU749750

Mont Ventoux 11 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 14Jul.2011 44.15158 5.32319 1466 Rumsais Blatrix 4 1 KU749649 KU749751

Mont Ventoux 12 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 14Jul.2011 44.15231 5.31939 1425 Rumsais Blatrix 4 KU749650 KU749752

Mont Ventoux 13 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 14Jul.2011 44.15183 5.32328 1466 Rumsais Blatrix 4 KU749651 KU749753

Mont Ventoux 14 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 14Jul.2011 44.1521 5.3232 1473 Philippe Wegnez KU749754

Mont Ventoux 15 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France Aug.2011 44.1521 5.3232 1473 Claude Lebas 4 KU749755

Mont Ventoux 16 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France Aug.2011 44.1521 5.3232 1473 Claude Lebas 4

Mont Ventoux 2 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 2009 44.15158 5.32319 1466 Christophe Galkowski 8 1 KU749653 KU749757 KU749792 KU749827

Mont Ventoux 3 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 2009 44.15158 5.32319 1466 Christophe Galkowski 3

Mont Ventoux 4 *Proformica longipilosa* sp. nov. Mont Ventoux, Chalet Reynard Bedoin (84) France 2009 44.15158 5.32319 1466 Christophe Galkowski 3

Vinsobre 1 *Proformica longipilosa* sp. nov. Vinsobre Vinsobre (84) France Aug.2012 44.33333 5.05972 297 Philippe Wegnez KU749654 KU749758 KU749793 KU749828

Appendix 2. Measurements of morphological characters for 321 workers of *Proformica* from southern France.

colony CW nCH nCU nTx nSc nG PDG GHL

Aurons 1 1190 1 2 2 2 5 17 70	Bonnieux 2 715 0 0 0 0 2 21 42
Aurons 1 850 0 2 1 0 5 18 50	Plateau de Calern 10 1020 7 8 20 3 18 42 155
Aurons 1 545 0 1 1 1 1 11 35	Plateau de Calern 10 900 5 6 23 4 10 42 136
Aurons 1 580 0 0 0 0 0 14 0	Plateau de Calern 10 731 7 2 10 3 5 42 102
Beaucaire 1 1240 0 1 0 0 2 14 42	Plateau de Calern 10 663 5 3 6 4 7 42 145
Beaucaire 1 1150 0 2 0 0 1 14 60	Plateau de Calern 18 1530 10 13 75 22 38 56 221
Beaucaire 1 990 0 1 0 0 2 14 50	Plateau de Calern 18 1275 8 8 37 11 28 56 187
Beaucaire 1 715 0 0 0 0 2 15 50	Plateau de Calern 18 1225 6 5 33 7 22 68 204
Beaucaire 2 815 0 0 0 0 1 17 60	Plateau de Calern 18 612 1 2 7 4 9 42 85
Beaucaire 2 600 0 0 0 0 0 20 0	Plateau de Calern 7 1070 6 8 27 4 28 56 170
Beaucaire 2 650 0 0 0 0 2 15 50	Plateau de Calern 7 663 6 5 13 2 10 42 155
Beaucaire 3 885 0 0 0 0 2 17 35	Plateau de Calern 2 1100 7 15 34 8 20 42 200
Beaucaire 3 595 0 0 0 0 0 11 0	Plateau de Calern 2 715 4 5 11 2 15 38 100
Beaucaire 3 615 0 0 1 0 2 15 50	Plateau de Calern 2 680 2 4 11 3 10 38 100
Beaucaire 4 935 0 1 0 0 3 14 50	Plateau de Calern 3 1200 11 10 27 8 39 38 220
Beaucaire 4 920 0 0 0 0 2 13 50	Plateau de Calern 3 835 9 7 21 5 27 42 170
Beaucaire 4 680 0 0 0 0 0 17 0	Plateau de Calern 3 715 2 8 12 6 16 42 170
Beaucaire 4 615 0 0 0 1 1 15 50	Plateau de Calern 4 1225 10 5 48 7 43 38 200
Bonnieux 1 600 0 0 0 0 0 17 0	Plateau de Calern 4 765 9 4 18 5 17 42 170
Bonnieux 1 680 0 0 0 0 2 18 35	Plateau de Calern 4 765 8 7 9 3 13 38 100
Bonnieux 1 700 0 0 1 0 0 21 0	Plateau de Calern 6 1395 13 17 65 8 36 42 200
Bonnieux 1 680 0 0 0 0 2 17 35	Plateau de Calern 6 700 8 5 21 4 15 38 100
Bonnieux 2 730 0 0 2 0 0 24 0	Plateau de Calern 6 735 4 5 18 5 11 42 170
Bonnieux 2 615 0 0 0 0 2 28 42	Plateau de Caussols 1 1350 8 6 45 9 26 56 190
Bonnieux 2 615 0 0 0 0 0 17 0	Plateau de Caussols 1 1170 9 8 40 5 22 48 170

Plateau de Caussols 1 715 6 4 18 2 13 42 102	Grospierres 10 815 2 1 3 1 4 10 50
Plateau de Caussols 19 1275 4 6 36 9 32 56 221	Grospierres 10 715 0 0 0 0 0 12 0
Plateau de Caussols 19 918 3 8 24 6 25 68 155	Grospierres 10 600 0 0 0 0 1 14 35
Plateau de Caussols 19 714 1 4 16 5 7 45 130	Grospierres 2 870 0 0 1 0 2 10 35
Plateau de Caussols 2 1325 12 9 57 10 38 48 170	Grospierres 2 615 0 0 1 0 0 12 0
Plateau de Caussols 2 1445 8 6 42 5 37 48 200	Grospierres 2 615 0 0 0 0 0 9 0
Plateau de Caussols 2 1120 6 9 30 5 29 48 170	Grospierres 3 850 0 0 0 0 2 9 35
Plateau de Caussols 3 1170 3 4 30 6 32 56 155	Grospierres 3 680 0 1 0 1 1 12 35
Plateau de Caussols 3 830 6 4 21 5 22 42 120	Grospierres 3 545 0 0 0 0 0 12 0
Plateau de Caussols 3 545 3 2 11 2 6 42 68	Grospierres 4 920 0 1 5 1 3 14 60
Plateau de Caussols 4 1260 6 7 44 7 32 56 190	Grospierres 4 715 0 0 0 0 0 12 0
Plateau de Caussols 4 730 4 5 14 4 12 42 102	Grospierres 4 650 0 0 0 0 1 10 35
Plateau de Caussols 5 1460 9 17 54 14 37 56 220	Grospierres 5 730 0 0 4 0 3 10 50
Plateau de Caussols 5 950 10 6 31 8 26 42 155	Grospierres 5 650 0 0 0 0 1 12 35
Plateau de Caussols 5 650 3 3 4 2 11 42 100	Grospierres 5 580 0 0 1 0 1 9 35
Plateau de Caussols 6 1325 15 13 60 12 35 56 190	Grospierres 6 920 1 0 5 0 4 10 50
Plateau de Caussols 6 1020 12 7 26 4 15 38 140	Grospierres 6 650 0 0 0 0 0 9 0
Plateau de Caussols 6 615 2 3 10 2 10 38 120	Grospierres 6 615 0 0 0 0 1 12 35
Plateau de Caussols 7 1360 10 11 46 7 31 48 170	Grospierres 7 935 1 0 6 4 3 10 35
Plateau de Caussols 7 780 5 3 12 4 18 42 100	Grospierres 7 750 0 0 3 0 2 9 35
Plateau de Caussols 7 680 3 3 9 3 8 42 100	Grospierres 7 665 0 0 1 0 1 12 35
Collias 1 1160 0 2 2 0 7 14 76	Grospierres 8 850 1 0 6 0 3 12 50
Collias 1 580 0 0 1 0 2 12 50	Grospierres 8 810 0 0 0 1 0 12 0
Collias 1 660 0 0 0 0 1 12 68	Grospierres 8 715 0 0 2 0 2 12 35
Collias 1 731 0 0 0 0 1 11 75	Grospierres 9 870 2 0 4 1 4 12 50
Collias 1 650 0 0 0 0 0 11 0	Grospierres 9 575 0 0 0 0 0 10 0
Plaine de la Crau 1 1000 0 2 7 0 2 18 50	Grospierres 9 550 0 0 0 0 0 10 0
Plaine de la Crau 1 870 0 0 0 0 2 14 35	Jonquières 1 1120 0 0 0 0 0 12 0
Plaine de la Crau 1 815 0 0 2 0 0 14 0	Jonquières 1 663 0 0 0 0 1 12 68
Plaine de la Crau 1 665 0 0 0 0 0 14 0	Jonquières 2 1050 0 0 0 0 2 14 50
Plaine de la Crau 1 615 0 0 0 0 2 12 45	Jonquières 2 714 0 0 0 0 1 12 68
Plaine de la Crau 2 885 0 3 3 0 1 15 60	Jonquières 3 952 0 0 1 0 1 14 60
Plaine de la Crau 2 850 0 2 5 2 0 17 0	Jonquières 3 1020 0 0 0 0 3 14 50
Plaine de la Crau 2 600 0 0 1 0 0 14 0	Jonquières 3 663 0 0 1 0 1 12 50
Plaine de la Crau 2 575 0 0 0 0 0 14 0	Grand Luberon 1 1395 6 5 28 3 10 14 85
Plaine de la Crau 2 650 0 0 0 0 1 14 50	Grand Luberon 1 1070 2 2 16 4 8 21 68
Plaine de la Crau 3 900 0 3 3 3 3 17 50	Grand Luberon 1 750 2 2 8 4 6 14 68
Plaine de la Crau 3 575 0 0 0 1 1 9 35	Grand Luberon 2 1000 4 5 14 5 8 14 68
Plaine de la Crau 3 615 0 0 0 0 0 11 0	Grand Luberon 2 970 4 6 16 1 6 21 68
Plaine de la Crau 3 715 0 1 0 0 1 11 35	Grand Luberon 2 665 4 1 8 1 3 21 50
Plaine de la Crau 3 600 0 1 0 1 2 12 50	Grand Luberon 2 580 2 0 5 0 2 24 50
Plaine de la Crau 4 1090 0 0 4 2 3 17 60	Grand Luberon 3 1290 7 4 21 2 7 17 85
Plaine de la Crau 4 800 0 0 0 0 1 14 35	Grand Luberon 3 800 3 4 10 3 7 34 68
Plaine de la Crau 4 850 0 0 1 0 3 14 60	Grand Luberon 3 680 4 3 4 3 1 34 35
Gréolières 7 1240 13 7 33 9 23 48 170	Grand Luberon 4 1020 3 4 13 2 3 14 68
Gréolières 7 850 8 4 11 5 15 42 155	Grand Luberon 4 665 4 1 3 4 5 24 85
Gréolières 7 665 2 3 10 3 10 42 100	Grand Luberon 4 630 4 2 5 3 5 34 60
Gréolières 8 1325 8 9 59 8 35 48 240	Grand Luberon 5 1200 8 6 16 4 14 14 68
Gréolières 8 815 6 7 17 5 9 42 155	Grand Luberon 5 935 4 0 3 1 7 34 68
Gréolières 8 645 3 4 12 3 7 38 140	Grand Luberon 5 760 4 3 6 2 7 34 60
Grospierres 1 850 0 2 3 2 1 10 50	Grand Luberon 6 1260 6 3 19 5 6 17 68
Grospierres 1 715 0 1 0 0 1 10 35	Grand Luberon 6 1020 2 3 12 5 9 17 68
Grospierres 1 560 0 0 0 0 0 10 0	Grand Luberon 6 665 3 2 7 1 3 14 50

Grand Luberon 7 1190 7 5 18 2 9 19 85	Saint-Baume 13 680 0 2 4 1 4 21 35
Grand Luberon 7 865 7 3 14 3 5 19 50	Saint-Baume 22 1080 2 4 16 5 8 15 60
Grand Luberon 7 700 2 0 0 0 0 34 0	Saint-Baume 22 865 1 3 11 1 7 15 50
Grand Luberon 8 1190 5 5 18 3 9 14 75	Saint-Baume 22 630 0 0 5 1 3 17 35
Grand Luberon 8 850 4 4 15 3 7 21 93	Saint-Baume 3 1190 2 3 19 3 3 20 50
Grand Luberon 8 650 4 0 6 0 4 28 50	Saint-Baume 3 833 0 5 8 3 2 17 50
Grand Luberon 9 1120 6 6 28 6 8 17 68	Saint-Baume 3 850 0 4 8 3 2 18 50
Grand Luberon 9 830 2 2 9 3 4 13 68	Saint-Baume 3 600 0 0 6 1 1 24 35
Grand Luberon 9 580 1 0 8 1 4 21 68	Saint-Baume 4 1140 6 5 12 3 6 15 70
Montagne de Lure 1 1225 6 4 12 7 10 14 68	Saint-Baume 4 1120 1 5 13 1 4 20 68
Montagne de Lure 1 765 2 1 9 4 8 14 50	Saint-Baume 4 1160 0 5 11 3 7 15 50
Montagne de Lure 1 700 0 1 5 0 7 14 42	Saint-Baume 4 645 0 0 2 1 2 12 50
Montagne de Lure 2 970 3 3 13 4 5 14 76	Saint-Baume 4 612 0 0 4 1 0 12 0
Montagne de Lure 2 815 1 1 7 2 9 14 50	Saint-Baume 5 1275 2 4 19 2 7 14 60
Montagne de Lure 2 700 1 2 7 3 5 14 50	Saint-Baume 5 1190 1 1 15 5 6 17 50
Montagne de Lure 3 1325 2 4 19 5 7 14 60	Saint-Baume 5 884 1 3 5 4 1 11 35
Montagne de Lure 3 815 2 3 11 1 6 14 50	Saint-Baume 5 665 0 3 5 0 2 14 50
Montagne de Lure 3 670 2 1 4 2 6 14 35	Saint-Baume 6 985 1 5 8 4 9 17 60
Montagne de Lure 4 1325 3 5 23 1 6 14 50	Saint-Baume 6 900 2 3 9 4 5 18 50
Montagne de Lure 4 850 5 5 9 2 12 14 65	Saint-Baume 6 630 2 2 6 2 5 14 50
Montagne de Lure 4 700 2 3 5 2 6 14 42	Saint-Baume 6 650 0 0 6 1 4 17 42
Montagne de Lure 5 1260 9 6 25 6 7 14 85	Saint-Baume 7 1050 3 3 9 4 10 13 50
Montagne de Lure 5 1020 3 2 15 3 9 12 50	Saint-Baume 7 870 2 2 7 3 6 17 50
Montagne de Lure 5 700 2 1 7 2 4 15 42	Saint-Baume 7 680 0 1 6 2 3 11 35
Montpellier 1 715 0 0 0 0 1 14 35	Saint-Baume 7 630 0 1 4 2 2 17 42
Montpellier 1 615 0 0 0 0 0 12 0	Saint-Baume 8 1155 4 4 17 4 10 14 50
Montpellier 2 630 0 0 0 0 1 17 35	Saint-Baume 8 935 2 4 7 2 6 15 42
Orange 1 715 0 0 11 2 7 48 120	Saint-Baume 8 800 3 3 5 2 5 17 68
Orange 1 884 1 0 5 1 8 42 140	Saint-Baume 8 665 0 1 5 3 3 14 35
Orange 1 680 1 2 10 0 4 42 120	Saint-Baume 9 730 2 4 9 2 7 17 50
Orange 1 715 2 1 9 1 5 42 135	Saint-Baume 9 815 1 4 8 3 8 14 60
Orange 1 1350 0 1 28 2 16 37 120	Saint-Baume 9 730 0 3 6 3 5 17 76
Orange 1 1140 0 2 24 4 19 42 120	Saint-Baume 9 650 0 3 6 3 5 14 50
Orange 1 865 0 1 12 3 8 56 135	Pompignan 1 1200 0 0 0 0 1 12 76
Orange 1 900 1 2 13 3 7 42 120	Pompignan 2 1170 0 2 2 0 3 11 68
Orange 2 580 0 0 3 1 1 42 85	Pompignan 3 1090 0 0 2 0 1 11.7 60
Orange 3 935 2 2 15 2 9 42 130	Sauteyrargues 1 750 0 0 0 0 0 12 0
Orange 3 815 1 0 4 2 4 42 110	Sauteyrargues 1 730 0 0 0 0 0 12 0
Orange 3 700 1 2 10 1 6 37 95	Sauteyrargues 2 600 0 0 0 0 1 13 35
Orange 3 715 2 2 5 2 5 42 110	Sauteyrargues 2 560 0 0 0 0 0 11 0
Saint-Baume 1 1140 6 6 21 4 7 15 60	Sauteyrargues 2 730 0 0 0 0 2 12 50
Saint-Baume 1 1105 2 5 12 4 5 18 50	Sisteron 1 800 10 5 18 3 11 34 130
Saint-Baume 1 815 2 6 5 3 2 13 75	Sisteron 1 645 8 3 13 4 10 34 120
Saint-Baume 11 1200 1 4 23 4 12 34 70	Sisteron 1 700 10 5 15 4 11 34 102
Saint-Baume 11 1100 3 1 20 1 5 21 50	Sisteron 2 1240 10 8 41 5 24 38 155
Saint-Baume 11 630 1 3 6 1 5 34 50	Sisteron 2 950 6 6 21 6 17 34 145
Saint-Baume 11 650 1 1 3 1 4 17 50	Sisteron 2 750 8 5 11 2 6 34 68
Saint-Baume 12 780 2 3 5 0 6 37 68	Sisteron 3 1035 10 9 35 6 18 34 160
Saint-Baume 12 680 2 1 8 2 4 17 42	Sisteron 3 800 10 3 13 4 10 42 102
Saint-Baume 12 730 1 3 2 1 1 14 35	Sisteron 3 680 6 2 12 4 16 34 110
Saint-Baume 12 650 2 0 7 3 2 17 50	Sisteron 4 815 10 7 20 4 16 34 155
Saint-Baume 13 985 4 2 7 2 7 14 35	Sisteron 4 750 6 4 11 2 8 34 95
Saint-Baume 13 730 0 3 2 1 4 17 25	Sisteron 4 665 8 3 14 1 9 42 85

Sumène 1 630 0 0 0 0 11 0	Mont Ventoux 12 1100 8 5 34 5 19 42 170
Sumène 1 665 0 0 0 0 11 0	Mont Ventoux 12 750 6 5 16 4 13 34 140
Sumène 1 560 0 0 0 0 11 0	Mont Ventoux 12 600 5 4 11 3 7 34 150
Sumène 2 1100 0 2 0 0 12 0	Mont Ventoux 13 1240 13 4 41 7 24 34 190
Sumène 2 615 0 1 0 0 11 0	Mont Ventoux 13 1190 14 6 33 5 12 37 170
Sumène 2 615 0 0 0 0 14 0	Mont Ventoux 13 730 5 4 16 5 7 42 140
Tarascon 1 1120 1 4 3 0 5 17 60	Mont Ventoux 13 665 5 3 14 4 7 42 140
Tarascon 1 920 0 2 1 0 2 17 50	Mont Ventoux 15 1300 13 5 38 9 22 28 205
Tarascon 1 920 0 2 1 0 2 14 45	Mont Ventoux 15 1150 10 2 37 10 16 34 190
Tarascon 1 715 0 0 2 0 2 12 50	Mont Ventoux 15 650 4 2 9 2 10 42 105
Tarascon 2 990 0 0 0 0 3 14 90	Mont Ventoux 15 680 1 0 11 3 9 37 85
Tarascon 2 870 0 0 0 0 2 15 50	Mont Ventoux 16 1020 8 4 32 7 25 42 190
Tarascon 2 650 0 0 0 0 12 0	Mont Ventoux 16 850 8 5 15 6 11 48 170
Tarascon 2 730 0 0 0 0 1 14 50	Mont Ventoux 16 640 8 3 11 3 9 48 155
Tarascon 3 1070 0 0 2 0 0 15 0	Mont Ventoux 16 650 6 2 12 4 9 48 170
Tarascon 3 715 0 0 0 0 1 14 50	Mont Ventoux 2 750 2 4 16 4 10 42 153
Tarascon 3 615 0 0 0 0 12 0	Mont Ventoux 2 697 2 3 11 5 10 42 153
Tarascon 3 615 0 0 0 0 2 13 50	Mont Ventoux 2 586 1 3 14 5 7 34 85
Mont Ventoux 1 1156 5 4 31 6 18 42 187	Mont Ventoux 2 1240 6 8 43 9 17 42 170
Mont Ventoux 1 646 3 2 11 7 13 42 170	Mont Ventoux 2 1326 9 6 46 12 17 42 195
Mont Ventoux 1 646 4 2 13 3 10 37 85	Mont Ventoux 2 1088 7 5 39 8 19 42 170
Mont Ventoux 10 1395 10 2 34 8 24 37 170	Mont Ventoux 2 697 2 4 17 3 10 34 93
Mont Ventoux 10 950 4 7 20 8 17 42 190	Mont Ventoux 2 730 3 4 19 4 12 42 136
Mont Ventoux 10 815 5 1 4 2 11 48 120	Mont Ventoux 3 1411 4 5 41 7 23 48 204
Mont Ventoux 10 665 4 4 9 1 7 37 85	Mont Ventoux 3 1122 7 0 30 7 16 48 187
Mont Ventoux 11 1310 9 6 47 8 26 42 170	Mont Ventoux 3 850 4 2 17 6 8 42 102
Mont Ventoux 11 850 6 5 18 5 11 48 170	Mont Ventoux 4 1326 8 9 44 9 21 56 187
Mont Ventoux 11 700 6 4 13 4 10 37 135	Mont Ventoux 4 1258 8 7 34 8 28 24 187
Mont Ventoux 11 665 3 2 13 5 7 37 170	Mont Ventoux 4 833 4 3 16 4 11 42 170
Mont Ventoux 12 1445 12 9 55 8 27 42 170	

Appendix 3. Measurements of morphological characters for 10 queens of *Proformica* from southern France.

colony	CW	SL	nCH	nCU	nPn	nMes	nPP	nSc	nG	PDG	GHL	MW	ML	ScW
Sainte-Baume	1410	1130	0	2	14	14	8	5	10	8	85	1070	2000	660
Tarascon 1	1500	1260	0	6	6	5	8	2	1	14	85	1090	2175	620
Beucaire 1	1530	1240	0	2	1	2	0	0	9	0	1020	2040	620	
Beucaire 1	1630	1275	0	4	0	0	2	0	2	11	70	1190	2245	665
Pompignan 10	1665	1235	1	2	1	10	6	2	3	11	70	1190	2245	800
Pompignan 10	1690	1240	0	2	1	11	5	2	2	11	85	1240	2310	865
Orange 1	1700	1275	3	2	11	26	10	2	25	15	240	1275	2340	765
Orange 1	1530	1225	1	3	15	30	6	3	24	17	210	1190	2245	700
Mont Ventoux 11	1770	1395	4	5	13	38	16	8	26	12	220	1410	2510	815
Mont Ventoux 2	1750	1325	13	7	20	53	15	13	36	14	240	1320	2450	850

Appendix 4. Relative proportions of cuticular hydrocarbons of *Proformica* workers. Hydrocarbons were identified using gas chromatography – mass spectrometry.

Peak no.	Species compounds	<i>P. longipilosa</i> (Mont Ventoux + Sisteron)		<i>P. nasuta</i> (Montagne de Lure + Grand Luberon)		<i>P. nasuta</i>		<i>P. longisetata</i>		<i>C. cursor</i>	
		mean	SE	mean	SE	mean	SE	-	mean	SE	
1	C23	0.82	0.37	0.23	0.14	0.00	0.00	2.29	1.34	0.21	0.00
2	11C23	0.29	0.12	0.06	0.05	0.00	0.00	0.00	0.00	0.00	0.00
3	7C23	0.35	0.12	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00
4	5C23	0.69	0.12	0.71	0.17	0.00	0.00	0.00	0.00	0.00	0.00
5	3C23	0.28	0.10	0.00	0.03	0.00	0.00	0.00	0.03	0.00	0.00
6	5,11+5,13C23	0.43	0.11	0.41	0.15	0.00	0.00	0.00	0.00	0.00	0.00
7	C24:1	0.00	0.00	0.00	0.00	0.00	0.00	4.41	0.00	0.00	0.00
8	C24	0.30	0.16	0.60	0.11	0.04	0.03	3.14	0.24	0.00	0.00
9	3,7+3,10C23	0.08	0.03	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
10	8+9+10+11+12C24	0.35	0.15	0.04	0.06	0.00	0.00	0.00	0.57	0.00	0.00
11	6C24	0.42	0.14	0.00	0.05	0.00	0.00	0.82	0.00	0.00	0.00
12	4C24	0.17	0.05	0.00	0.02	0.00	0.00	1.10	0.00	0.00	0.00
13	8,12C24	0.00	0.00	0.00	0.00	0.00	0.00	0.64	0.00	0.00	0.00
14	C25:1	0.21	0.10	0.00	0.03	0.02	0.02	0.00	0.00	0.00	0.00
15	6,8+6,10C24	0.16	0.06	0.02	0.07	0.03	0.03	0.00	0.00	0.00	0.00
16	C25	7.42	0.74	0.49	0.63	0.48	0.19	4.41	3.01	0.00	0.00
17	9+11+13C25	9.58	0.98	0.20	0.82	0.08	0.05	0.80	0.16	0.00	0.00
18	7C25	0.27	0.27	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00
19	5C25	10.16	1.03	0.16	0.88	0.00	0.00	1.51	0.00	0.00	0.00
20	9,13+9,15C25	0.17	0.06	0.32	0.08	0.09	0.06	0.53	0.00	0.00	0.00
21	3C25	1.73	0.37	0.16	0.17	0.02	0.02	0.30	0.26	0.00	0.00
22	7,15+7,17C25	0.15	0.15	0.01	0.04	0.05	0.04	0.00	0.00	0.00	0.00
23	5,9+5,13+5,15C25	13.37	1.78	0.20	1.19	0.00	0.00	0.00	0.00	0.00	0.00
24	C26:1	0.00	0.00	0.00	0.01	0.02	0.02	1.01	0.00	0.00	0.00
25	C26	0.36	0.10	0.64	0.11	0.02	0.01	2.92	0.37	0.00	0.00
26	3,7 C25 + 3,8+3,10C25	0.88	0.59	0.09	0.15	0.00	0.00	0.00	0.00	0.00	0.00


Peak no.	Species compounds	<i>P. longipilosa</i> (Mont Ventoux + Sisteron)		<i>P. nasuta</i> (Montagne de Lure + Grand Luberon)		<i>P. nasuta</i>		<i>P. longiseta</i>		<i>C. cursor</i>	
		mean	SE	mean	SE	mean	SE	-	mean	SE	
27	8+9+10+11C26	2.91	0.54	0.22	0.27	0.05	0.04	0.24	0.00	0.00	
27b	11,13,15+13,15,17TriMeC25	0.00	0.00	0.06	0.10	0.00	0.00	0.00	0.00	0.00	
28	8,12 DiMe C26	0.00	0.00	0.00	0.01	0.00	0.00	0.63	0.00	0.00	
29	6C26	0.79	0.14	0.04	0.08	0.05	0.03	0.55	0.00	0.00	
30	5C26	0.04	0.04	0.00	0.02	0.02	0.02	0.00	0.00	0.00	
31	4 Me C26	0.68	0.11	0.00	0.06	0.00	0.00	2.36	0.00	0.00	
32	11,13,15+13,15,17C25	0.00	0.00	0.00	0.00	0.00	0.00	0.24	0.00	0.00	
33	C27:1	0.00	0.00	0.00	0.01	0.01	0.01	0.00	0.00	0.00	
34	6,8+6,10C26	0.75	0.25	0.13	0.11	0.14	0.08	0.00	0.00	0.00	
35	4,8+4,10C26	1.79	0.47	1.30	0.27	0.03	0.03	0.00	0.00	0.00	
36	C27	2.58	0.43	1.97	0.31	0.45	0.14	7.14	2.98	0.00	
36b	4,8,12 TriMeC26	0.00	0.00	0.07	0.04	0.00	0.00	0.00	0.00	0.00	
37	9+11+13 Me C27	2.20	0.85	0.85	0.33	0.12	0.04	8.69	1.12	0.00	
38	7C27	0.12	0.10	0.35	0.07	0.07	0.06	1.05	0.00	0.00	
39	5 Me C27	4.83	0.79	3.01	0.49	0.03	0.02	1.70	0.72	0.00	
40	9,13+11,13C27	0.00	0.00	0.00	0.13	0.07	0.03	0.36	0.00	0.00	
41	3 Me C27	0.78	0.18	0.23	0.61	0.17	0.09	2.33	1.39	0.00	
42	5,9+5,15+5,17C27	25.22	3.15	4.30	2.15	0.02	0.01	0.72	0.62	0.00	
43	3,9C27	0.00	0.00	0.16	0.03	0.00	0.00	0.00	0.00	0.14	
44	C28	0.03	0.03	0.69	0.13	0.05	0.02	0.92	0.97	0.08	
45	5,9,13TriMeC27	0.00	0.00	0.04	0.04	0.00	0.00	0.00	0.00	0.00	
46	3,7+3,9C27	0.81	0.23	0.22	0.11	0.00	0.00	0.00	0.46	0.00	
47	8+9+10+12+14 Me C28	0.00	0.00	0.11	0.25	0.00	0.00	2.07	0.97	0.00	
48	13,15,17TriMeC27	0.00	0.00	0.05	0.10	0.00	0.00	0.00	0.00	0.00	
49	8,12+8,14+10,12+10,14DiMe C28	0.00	0.00	1.05	0.27	0.00	0.00	0.00	0.00	0.00	
50	7 Me C28	0.00	0.00	0.00	0.00	0.00	0.00	0.73	0.00	0.00	
51	6 Me C28	0.28	0.06	3.71	0.43	0.03	0.03	0.00	0.73	0.00	
52	4 Me C28	0.29	0.07	1.99	0.27	0.00	0.00	0.00	0.33	0.00	

Peak no.	Species compounds	<i>P. longipilosa</i> (Mont Ventoux + Sisteron)		<i>P. nasuta</i> (Montagne de Lure + Grand Luberon)		<i>P. nasuta</i>		<i>P. longiseta</i>		<i>C. cursor</i>	
		mean	SE	mean	SE	mean	SE	-	mean	SE	
53	9,13C28	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00
54	10,12+10,14 DiMeC28	0.00	0.00	0.22	0.27	0.03	0.02	1.84	0.00	0.00	0.00
55	7,11,19 TriMe C27	0.00	0.00	0.00	0.00	0.00	0.00	0.80	0.00	0.00	0.00
56	C29:1	0.00	0.00	0.00	0.07	0.02	0.02	0.00	0.00	0.00	0.00
57	6,10+6,12+6,14 DiMe C28	0.45	0.15	2.40	0.30	0.00	0.00	0.00	0.00	1.07	0.00
58	C29:1	0.00	0.00	0.00	0.15	0.25	0.10	0.00	0.00	0.00	0.00
59	4,8+4,10+4,12 DiMe C28	0.50	0.11	9.49	1.16	0.84	0.27	0.00	0.00	0.00	0.00
60	C29	0.31	0.10	5.16	0.60	1.41	0.34	2.57	0.00	6.60	0.00
61	4,8,11+4,8,14+4,10,14C28	0.02	0.02	6.05	1.29	0.05	0.05	0.00	0.00	0.00	0.00
62	9+11+13+15 Me C29	0.11	0.04	0.99	1.74	0.77	0.24	4.60	0.00	0.00	0.00
63	7 Me C29	0.48	0.45	0.26	0.25	0.85	0.23	6.37	10.72	0.39	0.89
64	5 Me C29	0.50	0.15	4.20	0.48	0.39	0.15	0.00	0.00	3.70	0.00
65	11,15+13,15 DiMe C29	0.00	0.00	0.02	0.79	0.17	0.06	3.49	0.00	0.00	0.00
66	9,13C29	0.00	0.00	0.00	0.27	0.00	0.00	0.00	0.00	0.00	0.00
67	3 Me C29	0.01	0.01	0.39	0.67	1.07	0.46	0.00	0.00	0.00	0.00
68	7,11+7,15+7,17 DiMe C29	0.05	0.05	0.00	0.01	0.00	0.00	16.41	19.53	0.00	0.00
69	5,9+5,15+5,17 Di Me C29	1.41	0.30	6.57	0.75	0.04	0.02	1.98	2.86	0.58	0.00
70	C30	0.13	0.06	0.00	0.29	1.44	0.33	0.00	0.00	0.00	0.00
71	3,9+3,11C29	0.00	0.00	0.14	0.08	0.00	0.00	0.00	0.00	4.54	0.00
72	9,13,15C29	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00
73	7,11,13+7,11,15C29	0.00	0.00	0.00	0.42	0.02	0.02	0.00	0.00	0.00	0.00
74	5,11,13+3,7,11C29	0.01	0.01	0.00	0.13	0.00	0.00	0.00	0.00	0.00	0.00
75	8+9+10+11+12 Me C30	0.00	0.00	7.04	0.84	0.87	0.38	1.27	1.30	0.16	0.00
76	11,13,15TriMeC29	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.00	0.00	0.00
77	3,11,13C29	0.01	0.01	0.02	0.18	0.01	0.01	0.00	0.00	0.00	0.00
78	8,12+8,14 DiMe C30	0.14	0.06	6.61	0.84	0.09	0.06	0.10	3.23	0.00	0.00
79	3+4 Me C30	0.00	0.00	0.63	0.25	0.29	0.10	0.00	0.00	0.00	0.00
80	9,11,17+9,13,15C29	0.10	0.05	1.14	0.23	0.00	0.00	1.74	0.00	0.00	0.00

Peak no.	Species compounds	<i>P. longipilosa</i> (Mont Ventoux + Sisteron)		<i>P. nasuta</i> (Montagne de Lure + Grand Luberon)		<i>P. nasuta</i>		<i>P. longiseta</i>		<i>C. cursor</i>	
		mean	SE	mean	SE	mean	SE	-	mean	SE	
81	6,10+6,12+6,14C30	0.00	0.00	0.82	0.21	0.09	0.04	0.00	0.00	0.00	0.00
82	8,12,16C30	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00
83	C31:1	0.01	0.01	0.00	0.05	0.03	0.02	0.00	0.00	0.92	0.00
84	4,8+4,10+4,12+4,14+4,16 DiMe C30	0.23	0.14	11.32	1.36	0.35	0.15	0.00	0.00	0.48	0.05
85	C31	0.23	0.07	0.12	0.32	1.58	0.39	0.54	1.52	0.03	0.03
86	8,12,13C30	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00
87	6,14,16+6,10,12+6,12,16C30	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00
88	4,10,12+4,12,16+4,14,16 TriMe C30	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00
89	9+11+13+15+16 Me C31	0.00	0.00	1.14	1.31	5.68	1.18	0.75	2.24	0.00	0.00
90	7C31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.84	0.00	0.00
91	13,15+13,17Me C31	0.00	0.00	0.00	0.65	0.04	0.04	0.00	0.00	0.00	0.00
92	11,13+11,15+11,17C31	0.00	0.00	0.00	0.21	0.00	0.00	0.00	0.00	0.00	0.00
93	9,11+9,13+9,15 DiMe C31	0.00	0.00	0.00	0.59	0.23	0.13	3.71	0.00	0.00	0.00
94	5C31	1.74	0.90	0.01	0.70	4.98	0.56	0.00	0.00	0.00	0.00
95	7,11+7,15C31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	13.26	0.00	0.00
96	5,9+5,13+5,15+5,17 DiMe C31	0.50	0.23	7.53	1.52	9.64	1.63	0.00	0.00	0.00	0.00
96b	11.15.19+13.15.19C31	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00
97	9,11,13+9,13,15+9,13,17C31	0.00	0.00	0.00	0.85	2.29	1.26	0.00	0.00	0.00	0.00
98	7,11,17+7,13,17C31	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.00
99	C32	0.00	0.00	0.44	0.58	3.11	0.67	0.00	1.24	0.00	0.00
100	3,11C31	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.00	0.00
101	5,10,14+5,10,12+7,11,15TriMeC31	0.17	0.07	0.30	0.54	1.95	0.74	0.00	0.00	0.00	0.00
102	8+10+11+12+13+14Me C32	0.01	0.01	0.65	0.43	2.81	0.45	0.00	1.14	0.00	0.00
103	11,13,15C31	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00
104	8,14+8,16+10,12+10,14DiMe C32	0.09	0.09	0.62	0.36	1.39	0.45	0.07	0.00	0.00	0.00
105	9,13,15C31	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00
106	6,10+6,14DiMe C32	0.00	0.00	0.32	0.13	0.81	0.12	0.00	1.02	0.00	0.00
107	4,12+4,14+4,16C32	0.00	0.00	0.08	0.05	0.00	0.00	0.00	0.00	0.00	0.00

Peak no.	Species compounds	<i>P. longipilosa</i> (Mont Ventoux + Sisteron)		<i>P. nasuta</i> (Montagne de Lure + Grand Luberon)		<i>P. nasuta</i>		<i>P. longisetata</i>		<i>C. cursor</i>	
		mean	SE	mean	SE	mean	SE	-	mean	SE	
108	C33:1	0.00	0.00	0.00	0.10	0.21	0.14	0.00	0.50	0.00	
109	C33	0.00	0.00	0.24	0.11	0.67	0.11	0.00	0.03	0.00	
110	4,12,xC32	0.00	0.00	0.00	0.13	0.50	0.18	0.00	0.00	0.00	
111	11+13+15+17 Me C33	0.02	0.02	0.04	0.47	3.03	0.45	0.00	0.55	0.02	
112	13,15 C33	0.03	0.03	0.05	1.07	5.35	1.26	0.15	0.00	0.00	
113	11,13+11,15C33	0.00	0.00	0.00	0.22	0.00	0.00	0.00	0.00	0.00	
114	5C33	0.00	0.00	0.00	0.42	1.00	0.58	0.00	0.00	0.00	
115	7,11+7,15C33	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.99	0.00	
116	5,11+5,13+5,15C33	0.02	0.02	0.03	1.12	8.47	0.68	0.00	0.14	0.08	
117	7,11,17C33	0.00	0.00	0.00	0.07	0.27	0.09	0.00	0.00	0.00	
118	C34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	
119	5,11,15+5,13,15C33	0.00	0.00	0.01	1.29	9.74	0.76	0.00	0.00	0.00	
120	10+12C34	0.00	0.00	0.00	1.86	9.89	2.10	0.00	0.14	0.04	
121	10,14+10,16+12,14C34	0.00	0.00	0.01	0.23	1.65	0.18	0.00	0.00	0.00	
122	8,12,16+8,12,18+8,14,16+8,14,16C34	0.00	0.00	0.00	0.17	0.94	0.19	0.00	0.00	0.00	
123	6,12+6,14+6,16C34	0.00	0.00	0.01	0.09	0.41	0.11	0.00	0.00	0.00	
124	C35	0.00	0.00	0.00	0.30	1.48	0.34	0.00	0.00	0.00	
125	11+13 C35	0.02	0.02	0.00	0.64	2.55	0.78	0.00	0.00	0.00	
126	11,15+13,15+13,17C35	0.00	0.00	0.00	0.30	1.35	0.37	0.00	0.00	0.00	
127	5,13+5,14+5,15+5,16C35	0.00	0.00	0.07	0.04	0.00	0.00	0.00	0.00	0.00	
128	C36	0.01	0.01	0.00	0.52	2.59	0.59	0.00	0.00	0.00	
129	5,13,15C35	0.00	0.00	0.00	0.26	1.56	0.24	0.00	0.00	0.00	
130	10+12+13+14+15+16+17+18C36	0.00	0.00	0.00	0.15	0.41	0.17	0.00	0.00	0.00	
131	12,16C36	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	
132	C37	0.00	0.00	0.00	0.16	0.85	0.18	0.00	0.00	0.00	
133	MeC37:1	0.00	0.00	0.00	0.10	0.34	0.12	0.00	0.00	0.00	
134	11+13+17 Me C37	0.00	0.00	0.00	0.07	0.10	0.09	0.00	0.00	0.00	
135	DiMeC37	0.00	0.00	0.00	0.20	0.79	0.27	0.00	0.00	0.00	
-	n =	8	16	22	1	5					

Appendix 5. Examples of chromatograms from cuticular extracts of *Proformica* workers analysed with gas chromatography.





DNA Library of Life

Corrigendum

The following corrections have been made to paper no. 290 (<https://doi.org/10.5852/ejt.2017.290>)

**Redescription of *Proformica nasuta* (Nylander, 1856)
(Hymenoptera, Formicidae) using an integrative approach –
Corrigendum**

Christophe GALKOWSKI¹, Claude LEBAS², Philippe WEGNEZ³,
Alain LENOIR⁴ & Rumsais BLATRIX^{5,*}

^{1,2,3,4,5} AntArea (www.antarea.fr), Association for the Study and
Mapping of Ants from Metropolitan France.

¹ 104 Route de Mounic, 33160 Saint-Aubin-de-Medoc, France.

² 2 Impasse del Ribas, 66680 Canohès, France.

³ Walbru (www.fourmiswalbru.com), Belgian Association for the Inventory of Ant Species in
Wallonia and Brussels, and Rue de la Grotte 23, 4651 Herve, Belgium.

⁴ IRBI, Institut de Recherche sur la Biologie de l’Insecte, UMR CNRS 7261,
Université François Rabelais, Faculté des Sciences, Parc Grandmont, 37200 Tours, France.

⁵ CEFE UMR 5175, CNRS – Université de Montpellier – Université Paul Valéry
Montpellier – EPHE, 1919 Route de Mende, 34293 Montpellier Cedex 5, France.

* Corresponding author: rumsais.blatrix@cefe.cnrs.fr

¹ E-mail: chris.gal@wanadoo.fr

² E-mail: cllebas@free.fr

³ E-mail: wegnez.phil@gmail.com

⁴ E-mail: alain.lenoir@univ-tours.fr

Galkowski C., Lebas C., Wegnez P., Lenoir A. & Blatrix R. 2017. Redescription of *Proformica nasuta* (Nylander, 1856) (Hymenoptera, Formicidae) using an integrative approach – Corrigendum. *European Journal of Taxonomy* 299: 1–2. <https://doi.org/10.5852/ejt.2017.299>

In the original publication, Fig. 1 mistakenly cited *Proformica pilinotum* sp. nov., which is a *nomen nudum*. The intended name was *Proformica longipilosa* sp. nov., which is reflected in the updated version of the figure provided below, which also includes a correction of the year of publication cited in the figure legend.




Fig. 1. Distribution of the 22 localities where *Proformica* nest samples were collected. The star indicates the type locality of *Proformica nasuta* (Nylander, 1856) (i.e., Beaucaire).

Published on: 14 March 2017

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d’Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands.